

Pig Age and Approachability Behavior to a Human Observer

A.S. Leaflet R3010

Shawna Weimer, Graduate Research Assistant;
Anna Johnson, Associate Professor;
Kenneth Stalder, Professor,
Department of Animal Science;
Locke Karriker, Associate Professor,
Veterinary Diagnostic and animal Production Medicine,
Iowa State University;
Thomas Fangman, Boehringer Ingelheim Vetmedica Inc.,
St Joseph, Missouri

Summary and Implications

The objective of this experiment was to determine if nursery pig age affected how many pigs touched and orientated to a human, or were classified as not-orientated during a human-animal interaction test using live human observation and digital image collection methodology. A complete randomized experimental design was utilized in this study where the pen of pigs was the experimental unit. Two methods, a human observer and a digital image, were assigned within rooms to all pens. Three age of pig treatments were compared; 6-weeks ($n = 6$ pens), 7-weeks ($n = 10$ pens) and 8-weeks ($n = 11$ pens). There were no differences for pig age when classified as touch, orientated and not orientated using the digital image ($P > 0.05$). Using the human method, more 7-week old pigs touched the human ($P = 0.02$). In conclusion, the digital image can be studied in more detail (on average each digital image was reviewed for 5 minutes) and hence, classifying pigs to these categories could be more accurate, resulting in no age differences for human approach in the pen between 6, 7 and 8 week old nursery pigs.

Introduction

Numerous human-animal tests in a variety of farm species have been used to try and measure fear levels. Such tests include the open field, human and novel approach-tests. Fangman and others (2010) coined the term “willingness to approach” (WTA) as a more positive alternative to fear. This WTA method allocated nursery pigs as either touching or orientating to the human in their home pen. A third category “not-orientated” included nursery pigs not meeting the previous criteria. The WTA method was conducted in real time by the human in the pen. In addition, how an animal reacts to a human can be vastly dependent upon the animals’ age and sex, as well as previous caretaker-pig interactions. Therefore, the objective of this experiment was to determine if nursery pig age affected how many pigs touched and orientated to the human or were classified as not-orientated during a human-animal interaction test using a live human observation and a digital image collection methodology.

Materials and Methods

Animal care and husbandry protocols for this experiment approved by the ISU-IACUC committee.

Location: The study was conducted November 4, 2010, at the Lauren Christian Swine Research Center at the Iowa State University Bilsland Memorial Farm, near Madrid, Iowa.

Animals: Purebred Duroc and Yorkshire crossbred barrows and gilts were used, body weight (BW) ranging from 24.4 kg to 31.9 kg, respectively. Pigs were not individually weighed before the study began. Average body weight was determined from previous performance records maintained on-site for nursery pigs of that age and genetic cross. All pens contained the same sex of nursery pig.

Housing and feeding: A total of 27 pens housed in three rooms were used. Each pen contained approximately 10 pigs per pen (0.32 m² per pig). Pens measured 1.5 m x 2.1m length, with steel dividers (81.3 cm height) between pens and one steel gate at the front of each pen (93.9 cm height.) A 4-hole dry feeder was located centrally at the front of the pen. Pigs were provided ad libitum access to a pelleted feed (1503 kcal/kg ME and 20.7% CP) formulated to meet or exceed requirements. Each pen contained one stainless steel nipple cup drinker 1.4 m from the front gate attached to the left or right pen divider, at a height of 33 cm above floor level. Metal tri-bar flooring was utilized in all pens. Caretakers observed all pigs at least once daily.

Experimental design: A complete randomized experimental design was utilized in this study where the pen of pigs was the experimental unit. Two methods, a human observer and a digital image, were assigned within rooms to all pens. Three age treatments were compared; 6-weeks ($n = 6$ pens), 7-weeks ($n = 10$ pens) and 8-weeks ($n = 11$ pens).

Approachability methodology: The methodology followed that previously described by Weimer et al (2014). On approach assessment day, a human observer approached the nursery pen, positioned the image-capturing device at the front of the gate at the approximate midpoint, and quietly stepped into the pen, immediately crouching down near the center of the gate. The evaluator extended and held still the left leather-gloved hand with the index finger extended, and began a stop watch, avoiding eye contact with the pigs for a 15-second period. The left hand and finger were extended to allow for the same anatomical location to be clearly visible in each digital image so that distance could be measured. At the end of the 15-second period, the observer looked behind her to ensure the sensor light on the digital camera had deployed and captured the digital image then looked back at the pigs and recoded the

live-categorical counts for the touch, orientated and not orientated categories. After counting all the nursery pigs, the observer rerouted her steps and exited the nursery pen. The live observation numbers for pigs' engaged in each of the three categories were recorded on a scan sheet located in the central alleyway. The observer then proceeded until all pens in the room had been entered, scanned and recorded. At the laboratory, each digital image was used to determine the three categories.

Measures: Live human observation and digital image were used to determine the number of pigs' engaged in touch, orientated and not orientated (Table 1). Pig percentages were calculated by dividing the total number of pigs classified in each category by the total number of pigs in the pen.

Table 1: Behavior classification of nursery pigs in a live human interaction test*

Measure	Description
Touch [1]	Any part of the pig's body touching the human observer
Oriented [2]	Pig oriented toward the human. Using Adobe Photoshop (Adobe Systems Incorporated, Arden Hills, Minnesota) in the digital image, a line was drawn from the midpoint between the pig's eyes to the center of the snout and then extended out towards the edge of the pen. If the line intersected with the human, the pig was classified as Orientated.
Not Oriented [3]	Pigs not exhibiting the above two behavioral classifications
	

Statistical Analysis: All data were evaluated for normal distribution before analysis by using the PROC UNIVARIATE procedure of SAS. A *P*-value of ≤ 0.05 was considered to be significant for all measures. Data were not normally distributed. These data were analyzed using the PROC GLIMMIX procedure of SAS. The main effect of pig age was compared (6-, 7-, and 8-weeks) for live and digital

image. The statistical model included the random effect of room. A Poisson distribution was noted for this data, hence the I-Link option was used to transform the mean and SE values back to the original units of measure for data and results interpretation.

Results and Discussion

There were no differences for pig age when classified as touch, orientated and not orientated using the digital image (*P* > 0.05; Table 2). Using the human method, more 7-week old pigs touched the human (Table 3; *P* = 0.02). In conclusion, the digital image can be studied in more detail (on average each digital image was reviewed for 5 minutes) and hence, classifying pigs to these categories could be more accurate, resulting in no age differences for human approach in the pen between 6, 7 and 8 week old nursery pigs.

Table 2: Least squares means (\pm SE) of average number and percentage of nursery pig age classified as Touch, Oriented, and Not Orientated by a digital-image

	Pig age (weeks)			SEM	P-value
	6	7	8		
Touch					
No. pigs/pen	2.5	2.3	1.7	0.5	0.52
Percent of pigs	26.4	24.5	18.3	4.1	0.30
Orientated					
No. pigs/pen	1.6	2.9	1.7	0.5	0.16
Percent of pigs	17.3	30.3	18.2	6.3	0.23
Not Orientated					
No. pigs/pen	5.3	4.3	6.0	0.7	0.26
Percent of pigs	56.3	45.2	63.5	7.0	0.16

Table 3: Least squares means (\pm SE) of average number and percentage of nursery pig age classified as Touch, Oriented, and Not Orientated by a human observer

	Pig age (weeks)			SEM	P-value
	6	7	8		
Touch					
No. pigs/pen	1.2 ^a	2.8 ^b	1.0 ^a	0.5	0.02
Percent of pigs	13.60 ^a	29.0 ^b	11.3 ^a	5.2	0.02
Oriented					
No. pigs/pen	2.4	3.4	2.0	0.6	0.19
Percent of pigs	24.4	35.2	21.8	5.9	0.18
Not Orientated					
No. pigs/pen	5.9 ^a	3.5 ^b	6.3 ^a	0.8	0.02
Percent of pigs	62.6 ^a	35.8 ^b	66.9 ^a	8.4	0.02

Acknowledgements

Boehringer Ingelheim Vetmedica Inc. for funding.