

Impact of 4R Management on Crop Production and Nitrate-Nitrogen Loss in Tile Drainage

RFR-A20103

Emily Waring, graduate student
Carl Pederson, agricultural specialist
Department of Agricultural and
Biosystems Engineering
John Sawyer, professor
Matthew Helmers, professor
Department of Agronomy
Terry Tuttle, farm superintendent

Introduction

Corn belt crop producers are increasingly challenged to maximize corn and soybean production while addressing the contributions farm practices make to the Gulf of Mexico hypoxia. Based on the need for nitrate-nitrogen (N) reductions to meet water quality goals, new management practices are needed to reduce nitrate-N losses at minimal cost and maximum economic benefits. This ongoing field research and demonstration project is evaluating various promising N management methods and technologies by documenting the nitrate-N export and crop yield from various systems. This is a summary of five years of crop yield data and four years of water quality data.

Materials and Methods

Funds provided by the Iowa State University Department of Agronomy Endowment helped to instrument the site for replicated studies of drainage water quality in 2013. In 2014, the site was uniformly cropped and the treatments were implemented for the 2015 growing season.

The site has 32 individual subsurface drained plots for drainage water quality evaluation. Drainage lines from individual plots are directed to separate sumps within culverts

where drainage is diverted for sampling. Each treatment is replicated four times. Treatments consist of corn-soybean rotation with each phase of the rotation present each year. The four treatments are:

1. Fall (anhydrous ammonia with nitrification inhibitor N-serve), 135 lb N/acre
2. Spring (anhydrous ammonia with no nitrification inhibitor), 135 lb N/acre
3. Split N, with 40 lb/acre of urea 2 x 2 starter at planting plus remainder in-season Agrotain treated urea, 135 lb N/acre total
4. No nitrogen applied

Tile flow water samples are analyzed for nitrate-N concentration. Additionally, the project documents crop yield for each treatment.

Results

Except for the early fall 2014 freezing conditions, which prevented fall anhydrous ammonia application (completed early spring 2015), agronomic operations were completed in a timely manner each year (Table 1). The 2015 year was characterized by greater than normal precipitation in late summer and fall, as well as a greater yearly precipitation than the 30-yr average (Cherokee, IA weather station about 10 miles south of the project site) (Table 2). The 2016 crop year also had more than the 30-yr average precipitation, with noticeably greater precipitation in April and September, which resulted in > 10 in. of average annual drainage (Table 3). The April precipitation delayed planting in 2016. The 2017 crop year had near normal precipitation in April and May, but much less than normal precipitation the rest of the year, resulting in < 4 in. of drainage on average. Precipitation in

2018 was above normal in May, June, and September with the total being 5 in. above normal for the year. The wettest year of the study was 2019 with 10 in. above normal precipitation, and most of the rainfall was from May to September. Precipitation in 2020 was well below normal after May.

In 2015, there was a 40 bushels/acre corn yield increase with N application in Treatments 1–3 as compared with Treatment 4 where no N was applied (Table 4). In 2016, the corn yield increase with N application was greater than 50 bushels/acre. During both 2015 and 2016, no statistically significant corn yield differences were observed between the treatments where N was applied. In 2017, corn yield increase with Treatments 1 and 2 compared with no N application was more than 75 bushels/acre. Also in 2017, there was a lower corn yield with the split N application compared with fall and spring ammonia timing. This was likely due to the limited precipitation after the sidedress N application and dry summer conditions (dry surface soil), which limited N movement into the soil and active corn root zone. In 2018, the split N application was timely with over an inch of precipitation a day after application. There was no statistically significant difference between the N application treatments in 2018, but all were significantly different compared with no N applied. In 2019 and 2020, there were no differences among N treatments, and they were all > 50 bushels/acre greater than the no N treatment. There were no statistical differences among the soybean yields in 2015,

which would be expected based on the uniform previous site history, no treatments applied to soybean, and no prior-year N applications to corn. Soybean yields in 2016 were greater than 70 bushels/acre for all treatments and greater than 60 bushels/acre in 2017. In 2018, soybean yield had very little variability in treatments and were 70–72 bushels/acre. Soybean yield for 2019 was not different among treatments, but all were between 58 and 63 bushels/acre. In 2020, soybean yield was significantly lower in the no N treatment.

Annual flow-weighted nitrate concentration was statistically similar whether N fertilizer was applied in the fall with inhibitor or in the spring in all years (Table 5). There was no N applied to soybean, but in 2016 (a very wet year) the concentrations with no N or split-N applied to the prior corn crop were still lower than with the fall or spring applied N, i.e. those treatment effects carried into the soybean year. Annual nitrate load was lowest in 2017 from all plots due to low drainage in the dry year (Table 6). There were not many differences among treatments in nitrate-N load due to high plot-to-plot flow variability.

Acknowledgements

Funds to conduct the research are being provided by the 4R Research Fund and the Foundation for Food and Agriculture Research.

Table 1. Dates of field operations for corn at the Northwest Research Farm.

	2015	2016	2017	2018	2019	2020
Fall NH ₃ + N-Serve application	4/18/2015	11/10/2015	11/16/2016	11/6/2017	11/16/2018	11/1/2019
Spring NH ₃ application	4/18/2015	4/12/2016	4/12/2017	4/29/2018	4/22/2019	4/1/2020
Planting date	5/18/2015	5/19/2016	5/6/2017	5/9/2018	5/7/2019	4/23/2020
Urea starter banded at planting	5/18/2015	5/19/2016	5/6/2017	5/9/2018	5/7/2019	4/23/2020
Agrotain treated urea sidedress	7/18/2015	7/14/2016	7/6/2017	7/3/2018	7/15/2019	7/8/2020
Harvest	10/18/2015	10/29/2016	10/24/2017	10/4/2018	10/16/2019	10/7/2020
Sulfur application		11/3/2016				
Planting population (seeds/acre)	34,000	34,000	35,077	35,077	35,077	35,077
Corn hybrid	Pioneer P0453	AgriGold 6267VT2RI B	Pioneer P0157AM X	Legend 9701	Wyfells W2506RIB	Wyfells W2506RIB

Table 2. Monthly precipitation from 2015 to 2020 at the Northwest Research Farm.

Mon	Precipitation (in.)						30-yr avg. precip. at Cherokee, IA (in.)
	2015	2016	2017	2018	2019	2020	
Jan	0.1	0.2	1.0	0.7	0.1	0.0	0.6
Feb	0.0	0.4	0.8	0.8	2.0	0.0	0.6
Mar	0.6	2.1	1.4	2.0	3.2	2.5	1.9
Apr	3.1	5.2	3.2	1.5	3.5	0.7	3.1
May	3.5	3.5	3.0	4.4	6.7	4.1	3.9
Jun	2.6	1.8	1.9	6.3	4.2	3.6	5.0
Jul	6.8	3.9	1.3	3.1	6.6	1.4	3.9
Aug	6.1	3.2	4.3	4.2	3.2	1.3	3.7
Sep	2.8	7.5	2.3	8.2	4.7	1.0	3.5
Oct	1.9	3.5	3.3	2.1	4.0	1.2	2.1
Nov	4.9	1.8	0.2	1.2	1.6	1.4	1.5
Dec	1.8	1.0	0.2	1.5	1.3	0.0	0.9
Total	34.1	34.0	22.9	36.0	41.0	17.1	30.7

Table 3. Drainage in inches by treatment from 2015 to 2020 at the Northwest Research Farm.

Nitrogen Management for Corn	Corn						Soybean					
	2015	2016	2017	2018	2019	2020	2015	2016	2017	2018	2019	2020
Fall NH ₃ (with inhibitor)	7.1	6.5	2.7	6.4	6.9	0.5	3.9	12.5	1.0	11.0	3.9	1.6
Spring NH ₃ (no inhibitor)	5.7	7.7	2.1	8.2	5.7	0.8	5.0	10.4	1.2	8.0	4.8	0.8
Split N	6.8	8.5	2.4	4.2	7.5	0.5	5.1	12.4	1.1	11.0	3.1	1.3
None	6.7	10.4	2.3	9.3	4.0	0.7	6.1	10.0	1.5	8.1	5.2	1.6

Table 4. Corn and soybean yields for 2015 - 2020 (bu/acre) at the Northwest Research Farm.

Nitrogen Management for Corn	Corn						Soybean					
	2015 ¹	2016	2017	2018	2019	2020	2015	2016	2017	2018	2019	2020
Fall NH ₃ (with inhibitor)	221a*	198a	203a	200a	214a	229a	62	74	62	70	58	69a
Spring NH ₃ (no inhibitor)	223a	200a	203a	195a	216a	231a	64	75	67	71	62	67a
Split N	224a	196a	181b	205a	207a	220a	64	72	66	70	59	66ab
None	183b	141b	125c	107b	137b	171b	61	74	64	72	63	59b

¹Early fall 2014 freezing conditions prevented fall anhydrous ammonia application (completed early spring 2015).

*Means with the same letter in the same column are not significantly different, P = 0.05. There were no significant differences among soybean yields.

Yields are reported at 15.5% moisture for corn and 13% moisture for soybean.

Table 5. Annual flow-weighted nitrate-N concentration (mg/L) from 2015 to 2020 at the Northwest Research Farm.

Year	Corn				Soybean			
	Fall NH ₃ (with inhibitor)	Spring NH ₃ (no inhibitor)	Split N	None	Fall NH ₃ (with inhibitor)	Spring NH ₃ (no inhibitor)	Split N	None
2015	16.1a*	15.7a	12.0ab	9.1b	12.7a	13.4a	12.1a	12.5a
2016	12.7a	12.5a	10.2a	9.8a	13.2a	13.7a	10.8ab	8.7b
2017	13.2a	13.8a	9.7a	11.8a	8.9ab	12.2a	8.6ab	5.7b
2018	10.4a	8.9ab	8.1ab	6.8b	11.0a	11.1a	8.3ab	6.4b
2019	10.6a	10.3a	8.1a	9.2a	10.6a	9.1ab	10.5a	5.8b
2020	9.2a	9.2a	8.8a	7.4a	11.6a	11.0a	10.1a	8.3a

*Means with the same letter in the same row and crop are not significantly different, P = 0.05.

Table 6. Annual nitrate-N load (lb/acre) from 2015 to 2020 at the Northwest Research Farm.

Year	Corn				Soybean			
	Fall NH ₃ (with inhibitor)	Spring NH ₃ (no inhibitor)	Split N	None	Fall NH ₃ (with inhibitor)	Spring NH ₃ (no inhibitor)	Split N	None
2015	24.4a	18.4a	16.6a	14.4a	9.8a	13.9a	13.2a	16.4a
2016	16.4a	20.7a	19.0a	21.6a	34.8a	30.4a	28.7a	19.1a
2017	7.5a	5.1a	4.7a	4.9a	2.3a	3.0a	2.1a	1.8a
2018	12.3a	15.7a	7.8a	13.8a	28.4a	19.0a	20.2a	11.3a
2019	16.3a	11.2a	13.2a	6.3a	7.6a	9.0a	7.4a	6.4a
2020	1.4a	1.6a	0.8a	1.1a	3.5a	1.8	2.8a	2.1a

*Means with the same letter in the same row and crop are not significantly different, P = 0.05.