



# On-Farm Demonstration Trial: Crop Protection Studies Approach® Fungicide Application on Corn

Mike Witt—on-farm trials coordinator and agronomist, ISU Extension and Outreach

Andrew Weaver—agricultural specialist, Northwest Research and Demonstration Farm

Dordt University, Agriculture Department

## Objective

Determine the effects of foliar fungicide application timing on corn yields to define best management practices.

## Introduction

An application of foliar fungicide to corn and soybean has become a common practice for many farmers in Iowa. The effect of fungicide on corn and soybean yield, however, can vary from year to year. Environmental conditions, such as rainfall and temperature, influence disease development, which will determine whether a fungicide affects yield. Because environmental conditions vary from one year to the next, it is difficult to predict how and when to use a fungicide. The objective of this trial was to evaluate whether the application of the foliar fungicide Approach® from Corteva corporation would result in a significant yield difference.

## Materials and Methods

### Crop Year–2021

Trial	210108
Trial County	Sioux
Soil Type	8B, 31, 91, 91B, 133, 310B, 310B2, 310C2, 428B
Previous Crop	Soybean
Tillage	Conventional
Current Crop	Corn
Hybrid–Variety Number	P0075AM
Hybrid–Variety Company	Pioneer/Corteva
Row Spacing	30 in.
Seeding Rate	34,000/ac.
Planting Date	May 1
Harvest Date	October 6
Fungicide	Approach 9oz./ac.
Fungicide Application	July 21
Experimental Type	On-Farm Demo
Replications	4

## Results



Trial Number	Treatment	Yield (bu./ac.) <sup>a</sup>	P-value <sup>b</sup>	Moisture	P-value <sup>b</sup>	Return on Treatment <sup>c</sup>
210108	Approach® 9oz./ac.	244.5 a	0.71	18.0 a	0.48	\$1077.59/ac.
	Control	243.3 a		17.8 a		\$1102.15/ac.

<sup>a</sup>Values denoted with the same letter within a trial are not statistically different at the significance level of 0.10.

<sup>b</sup>P-value = the calculated probability that the difference in yields can be attributed to the treatments and no other factors. For example, if a trial has a P-value of 0.10, there is 90% confidence the yield differences are in response to treatments. This is consistent for demonstration trials.

<sup>c</sup>Profit of Production based on \$17/ac. cost of Veltyma® product and \$12/ac. application cost, \$4.53 corn commodity prices. ((Yield x Price)-Costs) Commodity price is the 2020 national average cash price for corn.

## Location Climate Analysis



## Key Takeaways

- The usage of Approach<sup>®</sup> fungicide did not show any significant yield difference between the untreated control and with the given application timings.
- The fungicide had no significant effect on grain moisture in the trial.
- Return on treatment was greater if no fungicide was applied given the parameters of calculation.

NOTE: The results presented are from replicated demonstration trials. Statistics are used to detect differences at a location and should not be interpreted beyond the single location.