

2013

Effects of Seed Treatments and a Soil-applied Nematicide on Corn Yields and Nematode Population Densities

Gregory L. Tylka

Iowa State University, gltylka@iastate.edu

Mychele Batista da Silva

Iowa State University, mychelebs@yahoo.com.br

Joel L. DeJong

Iowa State University, jldejong@iastate.edu

Joshua L. Sievers

Iowa State University, sieversj@iastate.edu

Ryan Rusk

Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Microbiology Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Tylka, Gregory L.; Batista da Silva, Mychele; DeJong, Joel L.; Sievers, Joshua L.; and Rusk, Ryan, "Effects of Seed Treatments and a Soil-applied Nematicide on Corn Yields and Nematode Population Densities" (2013). *Iowa State Research Farm Progress Reports*. 1969. http://lib.dr.iastate.edu/farms_reports/1969

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Effects of Seed Treatments and a Soil-applied Nematicide on Corn Yields and Nematode Population Densities

Abstract

Plant-parasitic nematodes are microscopic worms that feed on plants. Almost every nematode that feeds on corn is capable of feeding on many other plants. These nematode parasites are thought to be native to most Iowa soils and to have fed upon native plants before corn was grown as a cultivated crop. Population densities (numbers) of most species of plant-parasitic nematodes that feed on corn have to increase to damaging levels (called damage thresholds) before yield loss occurs. Products that are currently available to manage plant-parasitic nematodes on corn in the state include the soil-applied insecticide/nematicide Counter® and two relatively new protectant seed treatments, Avicta® and Votivo®.

Keywords

Plant Pathology and Microbiology, Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Microbiology | Plant Pathology

Effects of Seed Treatments and a Soil-applied Nematicide on Corn Yields and Nematode Population Densities

RFR-A1255

Gregory Tylka, professor
 Mychele Batista da Silva, graduate student
 Department of Plant Pathology and
 Microbiology
 Joel DeJong, extension field agronomist
 Josh Sievers, farm superintendent
 Ryan Rusk, former farm superintendent

Introduction

Plant-parasitic nematodes are microscopic worms that feed on plants. Almost every nematode that feeds on corn is capable of feeding on many other plants. These nematode parasites are thought to be native to most Iowa soils and to have fed upon native plants before corn was grown as a cultivated crop.

Population densities (numbers) of most species of plant-parasitic nematodes that feed on corn have to increase to damaging levels (called damage thresholds) before yield loss occurs.

Products that are currently available to manage plant-parasitic nematodes on corn in the state include the soil-applied insecticide/nematicide Counter[®] and two relatively new protectant seed treatments, Avicta[®] and Votivo[®].

Counter[®] is a contact and systematic nematicide with the active ingredient terbufos. Avicta[®] is a contact nematicide with the active ingredient abamectin that moves on the surface of the root, and Votivo[®] is a special strain of the natural soil bacterium *Bacillus firmus* that grows on the root. Counter[®] is available from AMVAC, Avicta[®] from Syngenta Seedcare, and Votivo[®] from Bayer CropScience.

The objective of this experiment was to assess and compare the nematode population densities and yields of corn growing in plots with and without the seed-treatment nematode protectants and the soil-applied nematicide Counter[®].

Materials and Methods

The experiment was conducted on the ISU Northwest Research and Demonstration Farm. There were four replications of five different treatments. Plots consisted of eight rows, spaced 30 in. apart that were 94 ft long. The experiment was planted on April 26 and harvested on October 3, 2012. Soil samples for nematode analyses were collected on April 30 and then again, with root samples, on June 11, 2012, when the corn crop was at the V6 growth stage. Soil samples consisted of 20 one-inch-diameter cores that were 12 in. deep collected from under the seed row of the center four rows of each plot. The nematodes were extracted from the soil, and root samples and plant-parasitic nematodes were identified to genus and counted. The treatments, all applied to a single lot of seed of a single corn hybrid, were:

1. Avicta[®] Complete Corn (which is Avicta[®] + Cruiser[®] + Maxim[®] Quattro)
2. Cruiser[®] + Maxim[®] Quattro
3. Counter[®] + Cruiser[®] + Maxim[®] Quattro
4. Poncho[®] (500) / VOTiVO[®] + Acceleron[®] fungicides
5. Poncho[®] 500 + Acceleron[®] fungicides

Treatments 1 and 2 varied only by the presence of Avicta[®] and are labeled “w/ Avicta[®]” and “w/o Avicta[®]” in Figure 1. Similarly, Treatments 2 and 3 varied only by the presence of Counter[®] and are labeled “w/ Counter[®]” and “w/o Counter[®]” and Treatments 4 and 5 varied only by the

presence of Votivo[®] and are labeled “w/ Votivo[®]” and “w/o Votivo[®].”

Results and Discussion

The primary plant-parasitic nematodes found in the experiment were the lesion (*Pratylenchus*) and spiral (*Helicotylenchus*) nematodes. Dagger (*Xiphinema*) and stunt (*Tylenchorhynchus*) nematodes also were found at very low numbers and were combined into the category of “miscellaneous.” Spiral nematode was the most numerous (Figure 1). At planting, there were no significant differences in numbers of individual nematode types or in total number of plant-parasitic nematodes among treatments.

Very few nematodes were recovered from the root samples collected when the corn was at the V6 growth stage, so those data were not included. In the soil samples collected at V6, there were fewer nematodes found in each sample than were present at the time of planting (Figure 1), which was unusual. There were no significant differences in numbers of individual nematode types or in total number

of plant-parasitic nematodes among treatments at V6.

The overall average yield of the corn in the experiment was 164.0 bushels/acre. Average yields for the five treatments ranged from 162.0 to 168.1 bushels/acre, with no significant differences in yield among treatments.

Summary

The nematode management products had no effect on corn yields in the experiment. The effectiveness of the nematode management products may be much more pronounced in fields with very damaging nematode species (like needle and sting nematodes) and in fields with much greater plant-parasitic nematode population densities.

Acknowledgements

We thank AMVAC for donating the Counter[®] and Bayer CropScience and Syngenta Seedcare for donating and applying the seed treatments used in this research. No endorsement is intended of products used in the experiment, nor criticism implied of products not included.




Figure 1. Mean population densities of plant-parasitic nematodes in soil samples at planting (“Pi”) and at the V6 corn growth stage for the five treatments studied in the experiment. All nematode numbers were less than the numbers needed to significantly reduce corn yields. “Misc.” = miscellaneous plant-parasitic nematodes present in very low numbers. There were no significant differences ($P > 0.10$) in numbers of nematodes among treatments for the samples collected near the time of planting (Pi) or when corn was at the V6 growth stage.