

2003

## Barley Variety Test

Ronald Skrdla  
*Iowa State University*

Jean-Luc Jannink  
*Iowa State University*

Follow this and additional works at: [http://lib.dr.iastate.edu/farms\\_reports](http://lib.dr.iastate.edu/farms_reports)



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Skrdla, Ronald and Jannink, Jean-Luc, "Barley Variety Test" (2003). *Iowa State Research Farm Progress Reports*. 1487.  
[http://lib.dr.iastate.edu/farms\\_reports/1487](http://lib.dr.iastate.edu/farms_reports/1487)

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Barley Variety Test

## **Abstract**

Sixteen varieties were included in the 2002 barley test at Nashua. Each variety was sown in three different plots to average the effects of soil variability. The varieties were planted March 28 at a rate of 2 bushels/acre. All barley plots were harvested on July 17.

## **Keywords**

Agronomy

## **Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences

## Barley Variety Test

Ron Skrdla, ag research specialist  
Jean-Luc Jannink, assistant professor  
Department of Agronomy

### Materials and Methods

Sixteen varieties were included in the 2002 barley test at Nashua. Each variety was sown in three different plots to average the effects of soil variability. The varieties were planted March 28 at a rate of 2 bushels/acre. All barley plots were harvested on July 17.

### Results and Discussion

Barley yields averaged 101.1 bushels/acre in 2002, which is 2.0 bushels/acre more than in 2001 (Table 1). Excel was the highest yielding line based on three years of data (2000–2002) whereas Conlon had the highest test weight across all locations for the lines that were tested during the three-year period.

Additional information on barley variety tests in the state can be found in the publication, "Iowa Crop Performance Tests—Oat and Barley, 1998–2002," which is available from county extension offices (Pm-1645) and at [www.public.iastate.edu/~jjannink/](http://www.public.iastate.edu/~jjannink/).

**Table 1. Performance of spring barley varieties tested at Nashua from 2000–2002.**

Variety	Yield <sup>1</sup>			3-yr Avg	Test weight <sup>2</sup> (lbs/bu)	Heading date <sup>3</sup> (June)	Plant height <sup>4</sup> (in.)	Straw yield <sup>5</sup> (T/A)
	2000	2001	2002					
Azure	74.5	104.0	97.4	91.9	48.9	36	36	2.3
Bowers	60.6	96.5	105.9	87.7	49.0	37	35	2.3
Chilten	50.5	96.3	85.1	77.3	49.8	36	37	2.1
Conlon	-	101.0	106.2	92.4	50.8	34	32	2.3
Drummond	-	95.8	102.5	88.0	50.0	36	32	2.0
Excel	77.6	110.4	112.9	100.3	50.3	38	32	2.0
Hazen	74.7	99.5	100.2	91.5	48.6	38	33	2.2
Kewaunee	74.5	110.0	104.2	96.2	48.2	36	34	2.2
Lacey	-	100.0	115.3	96.4	50.7	37	33	1.9
Legacy	-	101.3	109.9	94.4	48.7	38	34	1.9
PrimusII	69.2	90.8	102.3	87.4	50.7	32	36	2.0
Robust	72.0	95.5	107.3	91.6	50.4	37	36	2.1
Royal	56.7	97.6	96.4	83.5	49.2	38	27	1.9
Seebe	-	88.1	85.7	75.7	49.8	45	33	2.2
Stander	58.0	95.8	101.9	85.3	49.6	38	30	2.0
Vivar	-	103.0	83.8	82.2	46.1	40	30	2.1
Average	66.8	99.1	101.1	88.9	49.4	37	33	2.1
LSD(0.05) <sup>6</sup>	7.0	10.0	8.9	10.3	1.0	1	1	0.3

<sup>1</sup>Grain yields are based on 48lb/bu. test weight.

<sup>2</sup>Test Weight – average from three sites.

<sup>3</sup>Data collected at Ames only.

<sup>4</sup>Height – Measured at Ames.

<sup>5</sup>Straw Yield – average from three sites.

<sup>6</sup>LSD = Least significant difference. When entries differ by an amount equal to one LSD or more, they are considered to be in different classes with 95% certainty.