

2010

## Soil Moisture

Joel L. DeJong

*Iowa State University*, [jldejong@iastate.edu](mailto:jldejong@iastate.edu)

Follow this and additional works at: [http://lib.dr.iastate.edu/farms\\_reports](http://lib.dr.iastate.edu/farms_reports)



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

DeJong, Joel L., "Soil Moisture" (2010). *Iowa State Research Farm Progress Reports*. 422.  
[http://lib.dr.iastate.edu/farms\\_reports/422](http://lib.dr.iastate.edu/farms_reports/422)

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Soil Moisture

## **Abstract**

Soil moisture samples were taken at 11 sites in northwest Iowa during the first few days of November, 2009. Moisture samples were pulled at 1-ft increments down to a 5-ft depth. Samples were weighed, oven dried, and reweighed at the Northwest Research Farm, Sutherland, IA.

## **Keywords**

RFR A9038, Agronomy

## **Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences

## Soil Moisture

### RFR-A9038

Joel DeJong, extension field agronomist  
ISU Extension

Soil moisture samples were taken at 11 sites in northwest Iowa during the first few days of November, 2009. Moisture samples were pulled at 1-ft increments down to a 5-ft depth. Samples were weighed, oven dried, and reweighed at the Northwest Research Farm, Sutherland, IA. The moisture percentage was calculated from these data and then used to calculate the inches of plant available moisture in the soil. The data from these sites are listed in the following table. Long-term fall averages range from about 4.5 in. to

6.0 in. in the top 5 ft of soil, but averages for the last 15 years have been significantly higher. The maximum plant available moisture level for most of these soils is around 11 in. in the top 5 ft of soil. Not all of the normal sites from NW Iowa had moisture samples pulled this fall. Late summer and fall precipitation in many neighborhoods gave an indicator that soil moisture profiles would be nearly full for the fall, so some of these areas were not sampled. Most areas that reported lower rainfall were included in this data set. Typical spring rainfall amounts should bring all sites to a full soil moisture profile by the time crops will need soil moisture reserves in 2010.

**Table 1. Soil moisture available to plants in inches.**

Site	County	2009 crop	Plant available moisture (in.)
Calumet	O'Brien	soybean	9.4
Sanborn	O'Brien	soybean	8.7
Doon	Lyon	soybean	9.3
Sibley	Osceola	soybean	7.8
Ireton	Sioux	corn	10.1
Hinton	Plymouth	corn	9.5
Aurelia (North)	Cherokee	soybean	9.7
Marcus (South)	Cherokee	soybean	8.7
Lawton	Woodbury	corn	8.6