

2001

Roundup Ready and Conventional Soybean Yield Trial

Kenneth T. Pecinovsky
Iowa State University, kennethp@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Pecinovsky, Kenneth T., "Roundup Ready and Conventional Soybean Yield Trial" (2001). *Iowa State Research Farm Progress Reports*. 1783.

http://lib.dr.iastate.edu/farms_reports/1783

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Roundup Ready and Conventional Soybean Yield Trial

Abstract

A yield performance study was conducted at the ISU NE Research and Demonstration Farm to compare twelve Roundup Ready® soybean varieties, and eleven conventional soybean varieties, one of which is a soybean variety grown for the organic market. Roundup Ultra was applied on the Roundup Ready® varieties, and a conventional herbicide program was used on the commercial soybean varieties to evaluate weed control and grain yield.

Disciplines

Agricultural Science | Agriculture

Roundup Ready[®] and Conventional Soybean Yield Trial

Ken Pecinovsky, superintendent

Introduction

A yield performance study was conducted at the ISU NE Research and Demonstration Farm to compare twelve Roundup Ready[®] soybean varieties, and eleven conventional soybean varieties, one of which is a soybean variety grown for the organic market. Roundup Ultra was applied on the Roundup Ready[®] varieties, and a conventional herbicide program was used on the commercial soybean varieties to evaluate weed control and grain yield.

Materials

The soil consisted of a Clyde silty clay loam and Floyd and Kenyon loam with a pH of 6.75 and with 6.1% organic matter. 1999 soil tests reported 38 ppm P205 and 148 ppm K20. The experimental design was a randomized complete block with three replications, and plots were 15 by 73 ft. The 1998 and 1999 crop was corn. The study used a conventional tillage system—fall chisel plowing and spring field cultivating prior to planting. Soybean varieties were planted 1.5 inches deep on May 15 at 182,402 seeds/acre except organic soybeans which were planted at 175,000 seeds/acre in 30 inch rows. Roundup Ultra was sprayed postemergently on June 27 at 35 oz per acre to the Roundup Ready[®] soybean varieties. The conventional soybean varieties were sprayed with 12 oz per acre Fusion, 1.15 oz per acre Pursuit, 2 oz per acre Cobra, 0.125 oz per acre Pinnacle, 0.25% V/V Activator 90 (non-ionic surfactant), and 32 oz per acre 28%

nitrogen on June 27. Plots were machine harvested for yield on September 30.

Results

Soybean variety harvest moisture, yield at 13% moisture, and income per acre are shown in Table 1. Yields were significantly different due to variety and herbicide program, if you exclude the HP-204 soybeans used for the organic market. The organically grown ISU HP-204's yielded 21.5 and 23.2 bushels per acre less than the RR[®] soybean and conventional varieties, respectively, but gained a premium of up to \$15 per bushel if grown as organic certified. Net income for RR[®] and conventional soybean varieties only differed by \$1.48 per acre. Conventional soybeans yielded about 1.7 bushels per acre greater compared with the RR[®] soybeans. The dollar savings on herbicide cost for the Roundup Ultra program more than offset the added cost of the technology fee on the RR[®] seed. Herbicide cost was approximately \$15/acre more for the conventional soybeans because four herbicides were required to control the weed species. Weed control was excellent for both herbicide programs.

Acknowledgments

We would like to thank the following companies for their cooperation on this research project—American Cyanamid Company, Asgrow Seed Company; Crows Seed Company, Golden Seed Company, Kruger Seed Company, LG Seed Company, Merschman Seed Company, Monsanto Chemical Company, Novartis Seed Company, Stine Seed Company, Pioneer Hi-Bred International, Inc.

Table 1. Yields of twelve Roundup Ready® and eleven conventional soybean varieties.

Variety	RR®	H ₂ O	Yield@13% moisture	Income/Acre**
Crows 2200-1	Yes	10.9	63.8	\$79.01
Novartis S24-K4	Yes	11.5	63.5	\$77.65
Stine 2506-4	Yes	11.1	63.2	\$75.83
Asgrow 2301	Yes	10.9	63.0	\$75.19
Pioneer 92B71	Yes	11.5	62.7	\$73.62
Kruger 202+	Yes	10.8	62.4	\$72.14
Stine 2101-4	Yes	10.9	61.5	\$67.71
Stine 1991-4	Yes	11.0	61.0	\$65.22
Asgrow 2401	Yes	11.1	60.3	\$61.60
Golden Hrv X92356	Yes	11.4	60.2	\$61.15
Pioneer 92B05	Yes	11.2	60.1	\$60.46
Stine 1901-4	Yes	11.0	59.3	\$56.40
Kruger 2525+	No	11.3	65.7	\$81.58
Mershman Com V	No	11.6	65.5	\$80.50
Golden Hrv X2120	No	10.7	64.7	\$76.32
Stine 2490-1	No	11.3	64.4	\$75.24
Stine 2499-0	No	11.1	63.5	\$70.59
Kruger 2343+	No	11.4	63.5	\$70.44
Pioneer 9233	No	11.0	63.1	\$68.35
Stine 2500	No	11.1	62.7	\$66.54
Asgrow 2247	No	10.9	61.2	\$59.09
LG Seeds 6200	No	11.7	60.3	\$54.44
HP204	No (with herbicide)	11.5	46.4	\$51.71
HP204	No (no pesticide)	11.1	40.3	\$301.40
Variety Average - LSD (P<0.05)		0.44	2.9	
Average (non-RRS) (excluding HP204)		11.2a	63.5a	\$70.31
Average (RRS)		11.1b	61.8b	\$68.83
Herbicide Program - LSD (P<0.05)		0.13	0.84	

** = Conventional soybeans @ \$5/bu, organic-certified soybeans @ \$15/bu (20% clean out), HP-204 soybeans with herbicide @ \$6.80/bu (20% clean out), non-RRS seed @ \$17/ac, RR® seed @ \$25/ac, P/K fertility maintenance @ \$20/ac, conventional herbicide cost @ \$35/ac, RR® herbicide cost @ \$20/ac, field operations @ \$50/ac, land rent @ \$125/ac.