IOWA STATE UNIVERSITY Digital Repository

Iowa State Research Farm Progress Reports

2002

Determining the Mechanism of Yield Stability in Alfalfa

Joe L. Robinson *Iowa State College*

Heathcliffe Riday *Iowa State University*

E. Charles Brummer *Iowa State University*

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports Part of the <u>Agricultural Science Commons</u>, <u>Agriculture Commons</u>, and the <u>Agronomy and Crop</u> <u>Sciences Commons</u>

Recommended Citation

Robinson, Joe L.; Riday, Heathcliffe; and Brummer, E. Charles, "Determining the Mechanism of Yield Stability in Alfalfa" (2002). *Iowa State Research Farm Progress Reports*. 1697. http://lib.dr.iastate.edu/farms_reports/1697

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Determining the Mechanism of Yield Stability in Alfalfa

Abstract

Year-to-year stability of crop yields is important for farmers and, hence, is an important goal of plant breeding programs. Especially in perennial crops like alfalfa, farmers need to know that they can count on consistent yields over a period of three or more years. Alfalfa varieties are composed of a population of many genetically distinct plants (or genotypes), unlike corn hybrids or soybean lines, which are genetically uniform.

Keywords

Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences

Determining the Mechanism of Yield Stability in Alfalfa

Joe Robins, graduate assistant Heathcliffe Riday, graduate assistant E. Charles Brummer, associate professor Department of Agronomy

Introduction

Year-to-year stability of crop yields is important for farmers and, hence, is an important goal of plant breeding programs. Especially in perennial crops like alfalfa, farmers need to know that they can count on consistent yields over a period of three or more years. Alfalfa varieties are composed of a population of many genetically distinct plants (or genotypes), unlike corn hybrids or soybean lines, which are genetically uniform.

Our goal is to determine whether the entire population of plants or single plants within a population determine yield stability of alfalfa. If varietal stability is due to the stability of individual plants in the population, then breeders can select individual plants with stable performance under many environmental conditions to use in the development of stable varieties. Conversely, if yield stability of a variety results from the interaction of many different genotypes, each of which performs better under some conditions than others, then alternative methods for developing stable cultivars must be investigated.

Previous research has suggested that increased genetic diversity can lead to increased stability; therefore, we will evaluate alfalfa varieties developed from different numbers of parental genotypes. If diversity leads to stability, those varieties derived from larger numbers of parent plants should perform more reliably. The results of this research will help breeders determine the number of parents and their genetic diversity when developing new, stable varieties.

Materials and Methods

We selected 10 individual genotypes from each of 10 varieties (5454, Affinity, DK140, Enhancer, Innovator +Z, Jade, Stampede, Wetland, WL324, and Vernal) for this study. The number of parental clones from which these varieties were developed ranged from eight to 99. Stem cuttings of each genotype were made in the greenhouse to allow for replication of individual plants. The cuttings were planted in five environments in Iowa, including Castana, in August 2000. The plot design was a lattice with two replications, with three cuttings per genotype in each replication. Yield was measured on each plot by hand-harvesting three times in 2000 and will be repeated in 2001. Yield stability over years and locations will be calculated at the end of the experiment.

Results and Discussion

After the 2002 harvest season, we plan to compare the average yield of each entire population with the yields of its component genotypes. We also will attempt to find a correlation between the yield stability of the populations and their parental clone numbers.

Acknowledgment

We thank Wayne Roush for his help in establishing and, especially, watering this experiment.