

**PLEISTOCENE LAND SNAILS FROM THE TALUS OF KENTUCKY
AND TENNESSEE**

LESLIE HUBRICH

3235 - 23rd Ave., Meridian, Miss. 39303

In areas where there is no loess, talus deposits are often the only source of Pleistocene fossils. Such talus deposits are found only in limestone areas, and are exposed where road or railroad construction has opened them up. They are important in Kentucky and Tennessee in showing that such species as *Hendersonia occulta* (Say) and *Allogona profunda* (Say) were once widely distributed here although they are not known to be living here at the present time.

KENTUCKY

1. FRANKLIN CO.: 2 miles south of Frankfort.

- Polygyra plicata* (Say)
- Stenotrema stenotrema* (Pfeiffer)
- S. leai aliciae* (Pilsbry)
- Mesodon thyroidus* (Say)
- M. clausus* (Say)
- M. zaletus* (Binney)
- M. appressus appressus* (Say)
- M. rugeli* (Shuttleworth)
- Triodopsis tridentata* (Say)
- T. vulgata* Pilsbry
- T. denotata* (Férussac)
- T. albolabris* (Say)

- Allogona profunda* (Say)
- Haplotrema concavum* (Say)
- Glyphyalinia indentata* (Say)
- Mesomphix vulgatus* H. B. Baker
- M. friabilis* (Binney)
- Ventridens gularis* (Say)
- V. ligerus* (Say)
- Anguispira alternata* (Say)
- A. kochi kochi* (Pfeiffer)
- Discus patulus patulus* (Deshayes)
- Succinea ovalis* (Say)
- Hendersonia occulta* (Say)
- Pomatiopsis lapidaria* (Say)

TENNESSEE

2. CHEATHAM CO.: Cumberland River bluff, opposite Ashland City.

- Stenotrema stenotrema* (Pfeiffer)
- Mesodon appressus appressus* (Say)
- M. inflectus* (Say)
- Triodopsis obstricta* (Say)
- Allogona profunda* (Say)
- Mesomphix vulgatus* H. B. Baker ?
- M. capnodes* (Binney)
- Discus patulus edentulus* Hubricht
- Hendersonia occulta* (Say)

- Helicina orbiculata* (Say)
Pomatiopsis lapidaria (Say)
3. CLAY CO.: 4 miles southwest of Celina.
Stenotrema angellum Hubricht
S. stenotrema (Pfeiffer)
Mesodon elevatus (Say)
M. rugeli (Shuttleworth)
Allogona profunda (Say)
Discus patulus (Deshayes)
4. JACKSON CO.: 10 miles northeast of Gainesboro.
Polygyra troostiana Lea
Stenotrema angellum Hubricht
S. stenotrema (Pfeiffer)
S. hirsutum (Say)
Mesodon zaletus (Binney)
M. elevatus (Say)
M. appressus appressus (Say)
M. rugeli (Shuttleworth)
M. inflectus (Say)
Triodopsis complanata (Pilsbry)
T. vulgata Pilsbry
Allogona profunda (Say)
Haplotrema concavum (Say)
Euconulus chersinus chersinus (Say)
Glyphyalinia lewisiana (Clapp)
G. praecox (H. B. Baker)
G. cryptomphala (Clapp)
Mesomphix vulgatus H. B. Baker
Paravitrea capsella (Gould)
Gastrodonta interna (Say)
Ventridenthes theloides (Walker & Pilsbry)
Anguispira alternata (Say)
Discus patulus (Deshayes)
Helicodiscus notius notius Hubricht
H. parallelus (Say)
H. aldrichianus (Clapp)
H. punctatellus Morison
H. jacksoni Hubricht
Succinea sp.
Strobilops labyrinthica (Say)
Gastrocopta armifera (Say)
G. contracta (Say)
G. holzingeri (Sterki)
- Gastrocopta pentodon* (Say)
Vertigo gouldi gouldi (Binney),
Carychium exile H. C. Lea
C. nannodes Clapp
Hendersonia occulta (Say)
Helicina orbiculata (Say)
Pomatiopsis lapidaria (Say)
5. JACKSON CO.: 8.5 miles northeast of Gainesboro.
Polygyra troostiana (Say)
Stenotrema angellum Hubricht
Mesodon zaletus (Binney)
M. elevatus (Say)
M. appressus appressus (Say)
M. rugeli (Shuttleworth)
Triodopsis complanata (Pilsbry)
Allogona profunda (Say)
Mesomphix vulgatus H. B. Baker
Anguispira alternata (Say)
Discus patulus (Deshayes)
Pomatiopsis lapidaria (Say)
6. SMITH CO.: 1 mile south of Carthage.
Polygyra troostiana Lea
Stenotrema stenotrema (Pfeiffer)
Mesodon zaletus (Binney)
M. elevatus (Say)
M. appressus appressus (Say)
M. inflectus (Say)
Triodopsis tridentata (Say)
T. obstricta (Say)
Allogona profunda (Say)
Haplotrema concavum (Say)
Mesomphix vulgatus H. B. Baker ?
Ventridenthes gularis (Say)
Anguispira alternata (Say)
Hendersonia occulta (Say)
Helicina orbiculata (Say)
Pomatiopsis lapidaria (Say)

MANUSCRIPT RECEIVED AND ACCEPTED FOR
 PUBLICATION SEPTEMBER 17, 1964.