

REPRINTS OF RARE PAPERS ON MOLLUSCA

REVIEW OF OUR PRESENT KNOWLEDGE OF

THE MOLLUSCAN FAUNA OF MICHIGAN

BY BRYANT WALKER

*Read before the Michigan Academy of
Science, December 27, 1894*

THE THOS. SMITH PRESS, 40-42 LARNED
STREET WEST
DETROIT

1895.

(Page 3)

I.

Barring a few scattering descriptions by European naturalists of such species as were brought home by the early travellers in this country, the history of North American conchology may be said to have begun when Thomas Say, in 1817, wrote the article on 'Conchology' for the first American edition of Nicholson's Encyclopedia of Arts and Sciences. Philadelphia then, as now, was the centre of activity in this branch of science in the United States, and in the proceedings of the then newly organized Academy of Natural Science and a few other scientific and literary publications of that city, nearly all the conchological writings for the next twenty years are to be found.

Michigan, as such, had no distinctive name in those days, and was known only as a wilderness filled with swamps and savages and located somewhere in that still greater and more indefinite region called the northwest.

But as population increased and young blood from the New England states made itself felt in the new territory, there began a dawn of better things. And one of the first acts of

the first legislature of the new state of Michigan in 1837 was the establishment of a State Geological Survey with Douglas Houghton at its head as Geologist and Dr. Abram Sager as Zoologist. Dr. Sager, who in after years became so well known in the medical department of the State University, and who had already in 1836 supplied Conrad with material for his monograph of the *Unionidae*, entered with activity upon the duties of his position and in 1839 published the first paper upon Michigan conchology. It is simply a list of species, 76 in number, one of which was not identified. It is dated January 12th, 1839, and is to be found

(PAGE 4)

in the Documents of the House of Representatives for 1839 at page 410.

In 1859, the Legislature passed an Act entitled 'An Act to finish the Geological Survey of the State.' The late lamented Prof. Alexander Winchell was appointed State Geologist and Prof. Manly Miles, who is still with us, State Zoologist.

The first report bears date December 31st, 1860. It contains in addition to other faunal lists, a catalogue of 161 species of shells, two of which *Planorbis truncatus* and *Unio leprosus* are described as new.

In the years, which had intervened between the publication of these catalogues, in addition to such scientific activity as centered around the labors of Dr. Sager and Prof. Winchell at the University, a little band of active collectors residing at Grand Rapids had done much to develop the fauna of the western part of the state. Alfred O. Currier, John A. McNeil, W. H. DeCamp and L. H. Streng were the leaders.

Mr. Currier came to Grand Rapids in 1850 from Troy, N.Y., where he had become fascinated with the study of conchology from being associated with that eminent conchologist, the late Dr. Wesley Newcomb. He died in 1880 and his extensive collection became the property of the Kent Scientific Institute of Grand Rapids. He published in 1859 (?) a 'List of Shells Col-

lected in the Grand River (Mich.) Valley,' and in 1865 a 'Catalogue of the Mollusca of Grand Rapids, Michigan.' In 1867 he published descriptions of four supposed new species from this state in the American Journal of Conchology, III, p. 112. In 1868 he published as No. 1 of the Miscellaneous Publications of the Kent Scientific Institute, an elaborate 'Catalogue of the Shell-bearing Mollusca of Michigan.' This list was by far the most complete yet published and enumerated 171 species and 6 varieties.

Dr. DeCamp came to Grand Rapids in 1855. In the congenial company of Mr. Currier he turned aside from botanical and geological work, which had previously enlisted his attention, and from that time has devoted his leisure hours almost wholly to the study of our local mollusca. He has accumulated a large and valuable collection, and his time and specimens have always been at the service of his fellow collectors. In 1881, Dr. DeCamp, under the auspices of the Kent Scientific Institute, published a 'Catalogue of the Shell-bearing Mollusca of Michigan,' in which, in addition

(PAGE 5)

to the list of 221 species and 9 varieties, he figured and described three species named by Mr. Currier, but never formally described.

In 1856, Mr. John A. McNeil settled in Grand Rapids and became interested in the subject through Mr. Currier and Dr. DeCamp. He remained there as an active and indefatigable collector until 1870, when he left and made collections in Central and South America a specialty. He died some three years ago at Binghamp, N. Y.

Mr. Streng, who has been a resident of Grand Rapids since 1870, began to collect as far back as 1850, when a resident of Saugatuck and is still actively engaged in the pursuit of his favorite study.

I am not aware that either of these gentlemen have ever published anything upon their Michigan collections. But Prof. Miles acknowledges the assistance afforded him by Mr. McNeil in the preparation of his catalogue. And Mr. Anthony was also indebted to him for some of the material from which he described a number of Michigan species, and, indeed, named one of them after him. The writer has elsewhere had occasion to express his obligations

to Mr. Streng for much generous assistance in compiling his previous catalogues of the shells of the state.

In 1879, the writer published a 'Catalogue of the Shell-bearing Mollusca of Michigan' in the Journal of Conchology and in 1892 a second list in the Nautilus.

In addition to the two papers by Mr. Currier, already referred to, the following local lists have been published:

In 1872-3, Mr. Sidney I. Smith published 'A Sketch of the Invertebrate Fauna of Lake Superior.'

In 1876, Mr. C. E. Beecher and myself compiled for the Ann Arbor Scientific Association a list of the species found in that vicinity.

And in 1893, I published 'A List of the Shells of the Saginaw Valley,' based upon the collection of the late Dr. George A. Lathrop.

In addition to these papers, which are devoted entirely to the shells of the state, many scattering references to our fauna are to be found in the writings of nearly all the prominent conchologists of this country. A full list of these will be found in the bibliography appended to this paper. Among them, however, are some worthy of special mention.

In 1836, T. A. Conrad in his 'Monography of the Unionidae' published descriptions of three supposed new forms from Michigan, viz: *Unio ellipsiformis*, *U. Sageri* and

(PAGE 6)

U. gibbosus var. *perobliquus*. The first of these species has never been identified by subsequent collectors and is probably the same species described in 1845 by Lea as *Unio spatulatus*. Unfortunately some of the details of Conrad's description are such, owing probably to the imperfect character of his specimens, as to prevent the union of the two species.

In 1847, William Case described in the American Journal of Science, a remarkable species of *Planorbis*, brought by Captain Stanard from Northern Michigan, under the name of *P. multivolvis*. The exact locality was not given. According to Binney a single specimen presented by Dr. Gould to the Smithsonian Museum was the only one known to him. Sowerby, in the 'Conchologia Iconica,' figured another form of the same species. I have not had an opportunity of examining that publication to obtain further details in regard to the specimen figured, but

it would seem that at least one other example had reached England. For many years nothing further was known of the species. Kuster in his monograph of the *Limnaeidae*, published in 1886, copied Sowerby's figure, and after criticizing both figures severely, pronounced it, in his opinion, an abnormal form of the common *Planorbis campanulatus* Say. Finally, however, in 1887, Dr. M. L. Leach found the long lost species in great abundance in Marl Lake in Roscommon county. These specimens agree perfectly with the original figures, and while examples from other localities seem to connect the typical form with *P. campanulatus*, it is at least a well marked variety, and of great interest, not only from its peculiar form, but from its remarkable history.

The naturalists connected with the celebrated expedition of Prof. Louis Agassiz to Lake Superior in 1848 found some seven new species of mollusca, two of which *Limnaea lanceata* and *Physa vinosa* were described by Dr. A. A. Gould, and the remainder, *Sphaerium aureum*, *emarginatum*, *flavum* and *tenuis* and *Pisidium rotundatum* by Temple Prime. The locality of most of these forms is given simply as Lake Superior, but they have been included in the catalogues of both Currier and De Camp. Some of them have since been definitely determined to be inhabitants of the state, and the remainder probably will be also.

In 1857, Dr. Isaac Lea (Proc. Acad. Nat. Sci. Phil. IX p 84) described the *Anodonta modesta* from specimens found near Kalamazoo.

In 1865 and 1866, John G. Anthony in the American

(PAGE 7)

Journal of Conchology described twelve new species of *Anodonta* and one *Unio* from Michigan. Of these but one, the *Anodonta subgibbosa*, is considered by Dr. Lea in his last 'Synopsis' to be a valid species.

In 1865, George W. Tryon, Jr., described a new species of *Limnaea*, the *L. zebra*, from specimens from Michigan, Minnesota and Wisconsin. It is now considered to be a color variety of the *L. reflexa* Say. And in 1866, the same author described the *Succinea DeCampii* from specimens discovered near Marshall in this state.

The growth of our knowledge of the molluscan fauna of the state during the fifty-five years which have elapsed since the first catalogue was published is shown by the following synopsis of the number of species listed in catalogues of Sager (1839), Miles (1860), Currier (1868), De Camp (1881), and the present one (1894). In arranging it all those species, whose occurrence in the state is considered doubtful for reasons hereinafter given, and all synonyms and varieties have been eliminated.

SUMMARY

	Land.	F. W. Pulmonates.	F. W. Operculates.	Bivalves	Total
Sager, 1839,	22	10	6	30	68
Miles, 1860,	44	24	14	57	139
Currier, 1868,	44	36	12	57	149
DeCamp, 1881,	48	41	27	69	185
Walker, 1894,	71	49	34	96	250

There have been up to the present time 42 species and varieties listed from this state as new to science. They are as follows:

Polygyra palliata alba Currier, Mss.
Vertigo Morsei Sterki

Succinea DeCampi Tryon.

Limnaea stagnalis Sanctae-Mariae Walker.
reflexa scalaris Walker
palustris Michiganensis Walker.
intertexta Currier, Mss.
contracta Currier.

(PAGE 8)

- Physa Parkeri* Currier.
deformis Currier.
Aplexa Tryoni Currier.
Planorbis truncatus Miles.
multivolvis Case.
bicarinatus corrugatus Currier Mss
bicarinatus major Walker
costatus DeTar & Beecher, Mss.
Valvata striata Lewis
Campeloma decisa flava Currier, Mss.
decisa melanostoma Currier, Mss
gibba Currier.
Milesii Lea.
Goniobasis Milesii Lea.
Unio ellipsiformis Con.
Sageri Con.
perplexus perobliquus Con.
leprosus Miles.
opalinus Anth.
Margaritina rhombica Anth.
Anodonta inornata Anth.
McNeillii Anth.
opalina Anth.
flava Anth.
glandulosa Anth.
imbricata Anth.
irisans Anth.
pallida Anth.
subinflata Anth.
subangulata Anth.
subgibbosa Anth.
subcarinata Currier
Houghtonensis Currier.
Sphaerium flavum Prime.

Of these the following have never been described formally, but have appeared by name simply in the different catalogues:

- Polygyra palliata alba* Currier.
Limnaea intertexta Currier.
Planorbis bicarinatus corrugatus Currier.
Campeloma decisa flava Currier.
decisa melanostoma Currier.

Of the remainder the majority, by reason of the increase of our knowledge of the variability of the species, have

(PAGE 9)

been already reduced to either varieties or synonyms. These are

- Succinea DeCampii* Tryon = *S. ovalis* Gld,
Physa Parkeri Currier = *P. Lordi* Bd,
Aplexa Tryoni Currier = *A. hypnorum* L.
Valvata striata Lewis = *V. sincera* Say

- Campeloma gibba* Currier = *C. rufa* Hald.
Milesii Lea = *C. subsolidum* Anth.
Unio Sageri Con. = *U. rectus* Lam.
leprosus Miles = *U. rectus* Lam.
opalinus Anth. = *U. novi-eboraci* Lea
Margaritina rhombica Anth. = *A. edentula* Say
Anodonta inornata Anth. = *A. decora* Lea
McNeillii Anth. = *A. Footiana* Lea.
flava Anth. = *A. fragilis* Lam.
glandulosa Anth. = *A. fragilis* Lam.
imbricata Anth. = *A. fragilis* Lam.
irisans Anth. = *A. fragilis* Lam.
pallida Anth. = *A. fragilis* Lam.
subcarinata Currier = *A. fragilis* Lam.
subinflata Anth. = *A. Maryattana* Lea
Houghtonensis Currier = *A. Maryattana* Lea
subangulata Anth. = *A. ovata* Lea.
Unio perplexus perobliquus Con. = *U. sulcatus* Lea. [Added with rubber stamp in my copy of this paper. A. L.]

II.

According to W.G. Binney, the leading authority on North American land shells, all that part of the continent east of the Rocky Mountains and north of Mexico, forms a single zoological province known as the Eastern Province. This again is divided into three regions; Northern, Interior, and Southern. The Northern Region comprises British America and that part of the United States lying east of the Appalachian chain of mountains while the Interior Region extends from the north region south to the alluvial lands lying along the Gulf of Mexico. Roughly speaking, the dividing line between the Northern and Interior Regions west of the Appalachian chain is the political boundary between Canada and the United States. But practically there is no hard and fast dividing line and one region gradually merges into the other. In the region of the Great Lakes, however, it would seem probable that the limits are more sharply defined.

Of the thirty-two species given by Binney as characteristic of the Northern Region, eight are peculiar to Greenland and Alaska. Of the remaining twenty-four, nineteen have been

(PAGE 10)

found in Michigan. While of the sixty-nine species belonging to the Interior Region forty-eight are known to inhabit this state. That is of sixty-seven Michigan species, nineteen or a little less than one-third belong to the

northern fauna and forty-eight to the interior fauna. The addition of the four species not included in Binney's list would not perceptibly change the proportion. This is what would be naturally expected from the position of the state upon the northern border of the Interior

Region and its very considerable longitudinal extent.

The seventy one species included in the present catalogue are divided among eleven families and sixteen genera as follows:

Family.	Genus.	Sub-Genus	Section.	No. of Species.
Selenitidae	Selenites			1
Limacidae	Limax			1
Vitrinidae	Vitrina			1
Zonitidae	Zonites		Mesomphix	3
			Hyalina	11
			Conulus	1
			Gastrodonta	2
Tebennophoridae	Tebennophorus			1
Endodontidae	Pyramidula	Pyramidula s.s.	Planogyra	1
			Goniodiscus	2
		Patula		2
		Helicodiscus		1
	Punctum			1
Helicidae	Polygyra		Triodopsis	12
			Stenotrema	3
	Acanthinula		Zoögenites	1
	Vallonia			3
Pupidae	Strobilops			2
	Pupa	Pupilla		1
		Leucochila		1
		Columella		1
		Bifidaria	Privatula	1
			Albinula	2
			Vertigopsis	2
		Angustula		1
	Vertigo	Vertigo s.s.		6
Stenogyridae	Ferussacia			1
Succineidae	Succinea			4
Auriculidae	Carychium			2
Total				71

The fluviatile fauna of the state includes representatives of six families and sixteen genera of univalves and two families and five

genera of bivalves. Of the univalves two families and six genera are pulmoniferous and four families and ten

(PAGE 11)

genera are branchiferous and operculate. The following is a list of the families and genera with the number of species belonging to each:

Family.	Genus.	No. of Species.
Limnaeidae	Limnaea	18 s.
	Planorbis	10

	Segmentina	2
	Ancylus	5
Physidae	Physa	11
	Aplexa	3
Valvatidae	Valvata	2
Viviparidae	Cameloma	5
Hydrobiidae	Bythinia	1
	Bythinella	4
	Somatogyrus	1
	Annicola	5
	Lyogyrus	1
	Pomatiopsis	2
Strepomatidae	Pleurocera	5
	Goniobasis	8
Unionidae	Unio	42
	Margaritina	5
	Anondonta	18
Corbiculidae	Sphaerium	19
	Pisidium	12
	Total,	179

The characteristic features of this fauna are best recognized by a comparison with those of other states. I have selected for that purpose Maine, New York, Indiana and Alabama, and have compiled the following table from the latest catalogues to which I have access.

I The table of states, catalogues, and numbers of species follows immediately after this paragraph in the original. It has been placed below in this reprint because of its width. The remainder of the text on page 11 follows. Ed. I

From an examination of these figures it will be noticed:

1st:-That the total number of species increases rapidly as we proceed from east to west and thence toward the south,

STATE	CATALOGUE	A	B	C	D	E	F	G	H	I	J*
Maine	Morse, 1864	48	2	26	2	4	1	0	10	12	57
New York	Lewis, 1874	64	2	33	2	5	4	7	52	20	123
Michigan	Walker, 1894	71	0	49	2	14	5	13	65	31	179
Indiana	Call, 1893	47	0	12	1	2	9	20	95	7	146
Alabama	Lewis, 1876	77	1	20	0	5	17	302	255	7	608

* In the original, the column headings, represented above by letters, are set vertically above each column. Their text follows. Ed.

A Land species

C Limnaeidae, Physidae

E Rissoidae

G Strepomatidae

I Corbiculidae

B Auriculidae (except Carychium)

D Valvatidae

F Viviparidae

H Unionidae

J Total fluviatile species

(PAGE 12)

The apparent exception in regard to Indiana must, I think be attributed rather to the lack of knowledge in regard to the extent of its fauna than to its actual paucity.

2nd:-That this increase is to be found almost wholly in the aquatic forms. This is shown not only by a comparison of the faunas of the different states but by the ratio in each state. Thus in Maine the land and fluviatile species are nearly equal; in New York the proportion is about 1 to 2; in Michigan 1 to

2½; in Indiana 1 to 3 and in Alabama 1 to 8. So also, while the increase in land fauna in Michigan and Alabama is about one half more than that of Maine, in the aquatic species the increase in Michigan is about 3½ times and in Alabama nearly 6 times.

3rd:-That this increase varies greatly in different families and between the northern and southern states is wholly confined to three: the *Viviparidae*, *Strepomatidae* and *Unionidae*; the rest suffering a radical reduction.

It follows that while the *Viviparidae*, *Streptomatidae* and *Unionidae* are thus shown to be essentially southern in their distribution, the *Limnaeidae*, *Physidae*, *Valvatidae*, *Rissoidae* and *Corbiculidae* are equally characteristic of the northern states.

4th:-Another interesting fact is developed upon examination of the distribution of three genera belonging to the *Unionidae* in the states above mentioned they are found as follows:

	<i>Unio</i> .	<i>Margaritina</i> .	<i>Anodonta</i> .
Maine,	3	2	5
New York,	34	6	12
Michigan,	42	5	18
Indiana,	73	8	14
Alabama,	238	13	5

From this it appears that while the genus *Unio* has evidently its metropolis in the south where it is enormously developed, the *Margaritinae* vary but little, and the *Anodontae* reach their maximum in the north. Thus while in *Unio* the proportion of Michigan to Alabama is about 1 to 6, in *Anodonta* the rate is reversed and is almost as much as the other way, i. e., 4 to 1.

The prominent features of our Michigan fauna then are the relative predominance of the *Limnaeidae*, *Physidae*, *Rissoidae*, *Corbiculidae* and *Anodontae* and in comparison with eastern states a large increase in the species of every family except the *Valvatidae* and *Viviparidae*.

This result is what might be expected from the situation and physical characteristics of the state. Surrounded on

(PAGE 13)

three sides by the great lakes, and with more than 5,000 small lakes with innumerable rivers and streams flowing into and out of them, and lying far enough north to afford a congenial habitat to the cold-water loving *Lymnaeidae*, *Physidae*, *Rissoidae* and *Corbiculidae*, many of whose species are circumpolar, Michigan stands pre-eminent in the number of species belonging to these families found within her borders, and there is every reason to believe that future investigation will tend to increase rather than diminish the list.

III.

But while the general features of our fauna are well enough known to enable it to be said that the present list of species will not pro-

bably be very largely increased in the future, yet it must be confessed that our present knowledge is very fragmentary, and requires much to be done before the perfect monograph of our mollusca can be written.

Large areas of our state are, as yet, almost wholly unexplored. The upper peninsula is practically a *terra incognita* to the conchologist. The authority for many of the species catalogued rests solely upon their occurrence at a single locality. Nor can it be said that the limits of the range north and south, east and west, of a single one of our 250 species are definitely and exactly known. With a part of our fauna coming to us from the north and east, and another part from the south and west, the accurate knowledge of the range of the different species over the state would enable us to solve many interesting questions relative to the origin and distribution of our mollusca. As an example of this, it may be stated that from our present information the rivers and lakes tributary to Lake Michigan appear to have a richer fauna than those that flow toward the east. These species belong mostly to the *Streptomatidae* and *Unionidae*, the characteristic families of the Mississippi Valley fauna. If this is found to be true, it would be in accord with the theory of the geologists, that, toward the end of the glacial period the great lakes had their outlet to the south into the Mississippi Valley, and tend to show that during that period these forms made their way north into Lake Michigan, and thence into its tributaries, but for some reason were prevented from effecting a lodgment in the eastern drainage of the state. Another interesting fact in the same connection is the recent discovery in the drift, near Toronto, Canada, of several Mississippi Valley species not now extant in the lake region.

(PAGE 14)

The existence of these species in Canada, and their subsequent extinction may have an important bearing upon the theory of a mild interglacial period, preceded and followed by an advance of the ice. If the ice receded to the vicinity of Toronto, allowing these Mississippi species to attain to that region, the fact that they did not establish themselves there would be easily accounted for by the subsequent advance of the ice and the extinction of the colony. The final melting and disappearance of

the ice cap, being complicated by changes in the direction of the drainage, might not afford a second opportunity for the immigration of the species in question.' (Simpson).

The first step towards the successful accomplishment of this purpose must be in the encouragement of local field work. Every local catalogue is a direct contribution to science whose value is only limited by the accuracy and thoroughness of the work it represents. The instruction in natural science which is, of late, becoming more and more incorporated with our public school system, could with great profit both to teacher and pupil be turned in the direction of this practical work. There are few branches of natural history better adapted for this purpose than conchology. The material is everywhere abundant and of great variety. It is easily collected and easily prepared for the cabinet and when once in suitable condition requires no further anxiety from its fortunate possessor. Then too, the technical literature required for this work is not extensive, and can be easily obtained. With the exception of the *Unionidae*, the various monographs published by the Smithsonian Institution contain practically everything that is necessary for the work of the ordinary collector, and these can be obtained without any great expense.

Then again, with the exception of the larger species of land mollusks, practically nothing is known of the development, life history and anatomy of even our most common species. Here is a most fruitful field for original investigation, which, now for many years, has been waiting for some new Say to immortalize himself by the elucidation of its problems.

The work done in the past in this state has been necessarily fragmentary, and it must continue to be so in the future, until such time as, under the direction of some supervising body of recognized authority, individual effort can be systematically organized and directed upon some well defined plan, which shall include the whole state.

(Page 15)

It would be exceedingly desirable, if under the stimulus, which it is hoped will be given to scientific matters in Michigan, by the organization of this society some such work could be set on foot as has been so successfully done by the Conchological Society of Great Britain and Ireland.

That organization in 1883 began a record of all localities in the British Isles from which the various species of mollusca were authentically known. For this purpose a committee was appointed of competent members, whose duty it was to keep an accurate record of every species. No record was allowed unless the locality where found was vouched for by some member or correspondent of the society, and the actual specimens were before the committee so that there could be no question as to identification. Under this system, during the ensuing ten years, over 31,000 records were made. While, of course, the details might have to be modified to meet the peculiar requirements of the work here, some scheme for similar work in this state would undoubtedly do much to stimulate interest and extend our knowledge.

But above all other things, what is needed in Michigan, is a complete biological survey of the state, conducted upon scientific principles and by scientific men under state auspices. This would naturally involve the establishment of a great state museum containing the results of the survey and the accompanying laboratories required to work up the material thus obtained. That this will come some time, I, for one, have no doubt, and I believe that this society should shape its policy with this end in view, and with this purpose collectively and individually seek to accumulate material in every branch of science, so that when the survey is established, and the natural history of Michigan comes to be written, we may all have our share in making it worthy of the great state we are proud to call our home.

(PAGE 16)

(PAGE 16)

CATALOGUE

of the

MOLLUSCA OF MICHIGAN.¹

	S	M	C	D	W
1 Selenites concsvus Say	x	x	x	x	x
2 Limax campestris Binn.					x
3 Vitrina limpida Gld.					x
4 Zonites fuliginosus Griff.	x	x	x	x	x
5 ligerus Say	x	x	x	x	x
6 inornatus Say	x	x			
7 cellarius Mull.					x
8 nitidus Mull.		x	x	x	
9 hydrophila Ing.			x	x	2
9 arboreus Say	x	x	x	x	x
10 radiatulus Alder					x
10 electrinus Gld.		x	x	x	
10 viridulus Mke.					x
11 indentatus Say		x	x	x	x
12 limatulus Ward		x	x	x	
13 minusculus Binn.		x	x	x	x
14 milium Morse					x
15 Binneyanus Morse					x
16 ferreus Morse					x
17 exiguus Stimp.		x	x	x	
18 fulvus Drap.					x
18 chersinus Say		x	x	x	
19 suppressus Say					x3x
20 multidentatus Binn.		x	x	x	
21 Tebennophorus Carolinensis Bosc					x
22 Patula alternata Say		x	x	x	x
22 alternata alba Tryon					x
23 solitaria Say		x	x	x	x
24 Pyramidula perspectiva Say		x	x	x	x
25 striatella Anth.		x	x	x	x
25 striatella alba					x
26 asteriscus Morse					x
27 Helicodiscus lineatus Say		x	x	x	x
28 Acanthinula harpa Say					x
29 Punctum pygmaeum minutissimum Lea					x
30 Helix virgata DeCosta* 4					x
31 Polygyra Mitchelliana Lea					x
32 clausa Say* 5		x	x	x	
33 multilineata Say		x	x	x	x
33 multilineata alba					x
33 multilineata unicolor					x
34 thyroides Say		x	x	x	x
34 thyroides bucculenta Gld.					x
35 albolabris Say		x	x	x	x
35 albolabris dentata Say					x
35 albolabris rufa DeKay					x
35 albolabris maritima Pils.					x

albolabris Traversensis Leach

Mss 6.

36 exoleta Binn.					x	x	x	x
36 zaleta Say								x
37 elevata Say					x	x	x	x
38 profunda Say					x	x	x	x
38 profunda alba								x

(PAGE 17)

	S	M	C	D	W
39 Polygyra Sayii Binn.		x	x	x	x
40 hirsuta Say		x	x	x	x
41 monodon Rack.					x
41 monodon fraterna Say		x	x	x	2x
42 Leaii Ward					x
43 tridentata Say		x	x	x	x
44 fraudulentata Pils.					x
44 fallax Say		x	x	x	2x
45 inflecta Say		x	x	x	x
46 palliata Say		x	x	x	x
46 palliata alba Currier, Mss.					x
47 Vallonia pulchella Mull.		x			x
47 minuta Say					x
48 costata Muller					x
49 excentrica Sterki					x
50 Strobilops labyrinthica Say		x	x	x	x
51 virgo Pils.					x
52 Pupa muscorum L.					x
53 fallax Say					x
54 edentula simplex Gld.		x	x	x	x
55 corticaria Say					x
56 armifera Swy		x	x	x	x
57 contracta Say		x	x	x	x
58 curvidens Gld.					x
59 pentodon Say		x	x	x	x
60 milium Gld.					x
61 Vertigo decora Gld.* 7					x3
62 ovata Say		x	x		x
63 Gouldii Binn.		x	x	2x	x
64 Bollesiana Morse					x
65 ventricosa Morse					x
65 ventricosa elatoir Sterki					x
66 tridentata Wolf					x
67 Morsei Sterki					x
68 Ferrussacia subcylindrica L.					x
68 lubrica Mull.		x	x	2	
69 Succinea aurea Say					x3
70 avara Say		x	x	x	x
70 vermeta Say					x
71 campestris Say* 8		x	x		
72 obliqua Say		x	x	x	x
72 ovalis Say		x	x		
73 ovalis Gld.					x
73 ovalis DeCampii Tryon					x
73 ovalis Higginsii Bld?* 9					x
73 ovalis Peoriensis Wolf Mss.					x

74	<i>Carychium exiguum</i> Say	x x x x	105	<i>pomilia Showalteri</i> Lea	x x
75	<i>exile</i> H. C. Lea	x	106	<i>brevispira</i> Lea	x
76	<i>Limnaea stagnalis</i> L.	x x x x	107	<i>deformis</i> Currier	x x
	<i>appressa</i> Say	x x2	108	<i>integra</i> Hald.	x x x
	<i>stagnalis jugularis</i> Say	x x		<i>Niagarensis</i> Lea	x x
	<i>stagnalis Sanctae-Mariae</i> Walker	x	109	<i>Aplexa hypnorum</i> L.	x x x
77	<i>ampla</i> Migh.	x x x		<i>elongata</i> Say	x x x2
78	<i>decollata</i> Migh.	x x x		<i>hypnorum Tryoni</i> Currier	x x x
79	<i>megasoma</i> Say	x x x		<i>hypnorum glabra</i> DeKay	15
80	<i>reflexa</i> Say	x x x x	110	<i>distorta</i> Hald.	x
	<i>umbrosa</i> Say	x x	111	sp.	x
	<i>reflexa zebra</i> Tryon	x x x	112	<i>Planorbis lentus</i> Say* 16	x
	<i>reflexa exilis</i> Lea	x	113	<i>trivolis</i> Say	x x x x x
	<i>reflexa scalaris</i> Walker	x		<i>corpulentus</i> Say	x3
	<i>reflexa distortus</i> Rossm* 10	x		<i>regularis</i> Lea	x x
	<i>reflexa Kirtlandiana</i> Lea	x	114	<i>truncatus</i> Miles	x x x x
81	<i>desidiosa</i> Say	x x x x x	115	<i>campanulatus</i> Say	x x x x x
82	<i>emarginata</i> Say	x x x		<i>campanulatus minor</i> Dkr.	x x x
83	<i>catascopium</i> Say	x x x	116	<i>multivolvis</i> Case	x x x
84	<i>caperata</i> Say	x x x x	117	<i>bicarinatus</i> Swy	x x x x x
85	<i>Cubensis</i> Pfr.	x		<i>bicarinatus major</i> Walker	x
	<i>umbilicata</i> Ads.	x x2x3x		<i>bicarinatus corrugatus</i> Currier Mss.	x x x
86	<i>pallida</i> Ads.	x x x		<i>bicarinatus Aroostookensis</i> Pils.	x
87	<i>humilis</i> Say	x x x	118	<i>exacutus</i> Say	x x x x
	<i>modicellus</i> Say	x x2		<i>exacutus</i> Say	x
	(PAGE 18)	S M C D W	119	<i>exacutus rubellus</i> Sterki	x
88	<i>Limnaea palustris</i> Mull.	x x x	120	<i>albus</i> Mull.	x x
	<i>elodes</i> Say	x x x2	121	<i>parvus</i> Say	x x x x
	<i>fragilis</i> L.	x	122	<i>deflectus</i> Say	x x x x
	<i>intertexta</i> , Currier Mss 11	x x	123	<i>costatus</i> DeTar & Beecher Mss 17	
	<i>palustris Michiganensis</i> Wkr.	x	124	<i>Segmentina armigera</i> Say	x x x x
89	<i>lanceata</i> Gld.	x x x	125	<i>Wheatleyi</i> Lea	x
90	<i>columella</i> Say	x x x x x	126	<i>Ancylus rivularis</i> Say	x
91	<i>contracta</i> Currier	x x x	127	<i>fuscus</i> Ad.	x x x x
92	<i>galbana</i> Say?	x	128	<i>parallelus</i> Hald.	x x x x
93	<i>bulimoides</i> Lea* 12	x	129	<i>diaphanus</i> Hald.	x x
94	<i>gracilis</i> Jay	x x x x	130	<i>tardus</i> Say	x x
95	<i>Binneyi</i> Tryon* 13	x x3		<i>Valvata tricarinata</i> Say	x x x x x
96	<i>Haydeni</i> Lea* 14	x3		<i>tricarinata bicarinata</i> Lea	x x
97	<i>Traskii</i> Tryon* 14	x3		<i>tricarinata unicarinata</i> DeKay	x
98	<i>Physa Lordi</i> Bd.	x x		<i>tricarinata simplex</i> Gld.	x x
	<i>Parkeri</i> Currier	x x	131	<i>sincera</i> Say	x x x x x
99	<i>gyrina</i> Say	x x x		(PAGE 19)	
	<i>elliptica</i> Lea	x			
	<i>oleacea</i> Tryon	x x3			S M C D W
	<i>Febigeri</i> Lea	x3		<i>Valvata sincera striata</i> Lewis	x x
	<i>gyrina Hildrethiana</i> Lea	x x x		<i>sincera Lewisii</i> Currier	x x
100	<i>heterostropha</i> Say	x x x x x	132	<i>humeralis</i> Say* 18	x
101	<i>Sayii</i> Tappan	x	133	<i>Vivipara contectoides</i> W.G. Binney* 19	
	<i>Sayii Warreniana</i> Lea	x x x	134	<i>Campeloma ponderosa</i> Say* 20	x x x
102	<i>vinosa</i> Gld.	x x x x		<i>decisa</i> Say	x x x x x
103	<i>anatina</i> Lea	x		<i>decisa flava</i> Currier Mss.	x
104	<i>ancillaria</i> Say	x x x			

	<i>decisa melanostoma</i> Carrier		179	<i>complanatus</i> Sol.	x	x3x
	Mss.	x		<i>purpureus</i> Say	x	
	<i>decisa heterostropha</i> DeKay	x x	180	<i>cornutus</i> Bar.		x x x
136	<i>integra</i> Say	x x2x x	181	<i>cuneolus</i> Lea* 28		x3
137	<i>rufa</i> Hald.	x x x x	182	<i>donaciformis</i> Lea		x
	<i>rufa gibba</i> Carrier	x x x	183	<i>elegans</i> Lea	x x x x	
138	<i>obesa</i> Lewis	x x x	184	<i>ellipsis</i> Lea	x x x x	
139	<i>subsolida</i> Anth.	x x		<i>olivarius</i> Raf.	x	
	<i>subsolida Milesii</i> Lea	x	185	<i>ellipsiformis</i> Con. 29		
140	<i>Lioplax subcarinata</i> Say* 21	x	186	<i>fabalis</i> Lea		x
141	<i>Bythinia tentaculata</i> L.	x3x				
142	<i>Bythinella attenuata</i> Hald.	x x				
143	<i>tenuipes</i> Couper* 22	x				
144	<i>Binneyi</i> Tryon 23	x				
145	<i>Nicklinana</i> Lea	x3x				
146	<i>obtusa</i> Lea	x x	187	<i>lapillus</i> Say	x x	
147	<i>Somatogyrus isogonus</i> Say	x x x x		<i>gibbosus</i> Bar.	x x x x	
148	<i>Ammicola limosa</i> Say	x x	188	<i>dilatatus</i> Raf.	x	
	<i>porata</i> Say	x x	189	<i>glans</i> Lea	x x x x	
	<i>pallida</i> Hald.	x x x x		<i>gracilis</i> Bar.	x x x x	
149	<i>grana</i> Say	x x2x x	190	<i>fragilis</i> Raf.	x	
	<i>granosa</i> Say	x	191	<i>iris</i> Say* 30	x x x x	
150	<i>Cincinnatiensis</i> Anth.	x x x	192	<i>Kirtlandianus</i> Lea 31		
151	<i>decisa</i> Hald	x	193	<i>laevissimus</i> Lea	x x x	
152	<i>lustrica</i> Pils.	x x	194	<i>Leibii</i> Lea		x
153	<i>Lyogyrus pupoideus</i> Gld.	x x	195	<i>latecostatus</i> Lea* 32		x x
154	<i>Pomatiopsis lapidaria</i> Say	x x x x		<i>ligamentinus</i> Lam.	x x x	
155	<i>Cincinnatiensis</i> Say	x	196	<i>crassus</i> Say		x2x3
156	<i>Pleurocera subulare</i> Lea	x x x		<i>luteolus</i> Lam.	x x x x	
	<i>subulare intensum</i> Anth.	x x	197	<i>siliquoideus</i> Bar.	x	x3
157	<i>neglectum</i> Anth.	x x2x x	198	<i>multiradiatus</i> Lea	x x	x3x
158	<i>elevatum</i> Say	x x	199	<i>nasutus</i> Say	x x	x x
159	<i>labiatum</i> Lea	x	200	<i>negatus</i> Lea* 33		x
160	<i>pallidum</i> Lea	x		<i>novi-eboraci</i> Lea	x x x x	
161	<i>Goniobasis livescens</i> Mke	x x x x	201	<i>opalinus</i> Anth.	x x	
	<i>Niagarensis</i> Lea	x x2	202	<i>occidens</i> Lea	x x x x	
	<i>livescens cuspidatus</i> Anth.	x3x	203	<i>parvus</i> Bar.	x x x x	
162	<i>translucens</i> Anth.	x3x	204	<i>penitus</i> Con* 34	x x	
163	<i>brevispira</i> Anth.	x3x		<i>perplexus</i> Lea* 35	x	
164	<i>pulchella</i> Anth.	x x2x	205	<i>gibbosus</i> Raf.	x	
165	<i>depygis</i> Say	x x x3x		<i>phaseolus</i> Hild.	x x x x	
166	<i>Milesii</i> Lea	x x x	206	<i>fasciolaris</i> Raf.	x	
167	<i>gracilior</i> Anth.	x3	207	<i>plicatus</i> Les.	x x x x x	
168	<i>semi-carinata</i> Say	x		<i>pressus</i> Lea	x x x	
169	<i>Virginica</i> Gml.* 24	x x	208	<i>compressus</i> Lea	x	
170	<i>Unio alatus</i> Say	x x x x x	209	<i>pustulatus</i> Lea* 36		x
171	<i>anodontoides</i> Lea* 25	x3		<i>pustulosus</i> Lea		x
172	<i>asperimus</i> Lea	x x	210	<i>bullatus</i> Raf.	x x	
173	<i>borealis</i> Gray	x		<i>radiatus</i> Lam* 37		
174	<i>caelatus</i> Con* 26	x x	211	<i>distans</i> Anth.	x x	
175	<i>Canadensis</i> Lea	x	212	<i>Rangianus</i> Lea		x
176	<i>cariosus</i> Say* 27	x x		<i>rectus</i> Lam.	x x x x x	
177	<i>circulus</i> Lea	x x x x		<i>leprosus</i> Miles	x x x	
178	<i>coccineus</i> Hild.	x x x x	213	<i>Sageri</i> Con 38		
				<i>rubiginosus</i> Lea	x x x x x	

(PAGE 20)

S M C D W

214	Schoolcraftii Lea	x x x x	247	Pepiniana Lea* 46	x x2x3x
	<i>prasinus</i> Con.	x2	248	plana Lea	x x x x
215	spatulatus Lea	x x x x	249	salmonia Lea	x x x
216	subovatus Lea* 39	x x x	250	Schaefferiana Lea	x x x
217	subrotundus Lea* 40	x x	251	Simpsoniana Lea	x
218	sulcatus Lea	x	252	subcylindracea Lea	x x x x
	<i>perplexus perobliquus</i> Con. 41		253	subgibbosa Anth.	x x x
219	Tappanianus Lea* 42	x3	254	<i>Sphaerium simile</i> Say	x x
220	tenuissimus Lea	x x x x		<i>sulcatum</i> Lam.	x x x x
221	trigonus Lea	x x x x	255	aureum Prime	x3
222	triangularis Lea	x x x x x	256	solidulum Prime	x x x x
223	undulatus Bar.	x x x3x		<i>distorta</i> Prime	x2
224	ventricosus Bar.	x x x x x	257	striatinum Lam.	x x x x
225	verrucosus Bar.	x x x x	258	rhomboideum Say	x x x
	<i>tuberculatus</i> Raf.	x	259	fabale Prime 51	x x
226	Margaritina complanata Bar.	x x x	260	occidentale Prime	x x x x
227	deltoides Lea	x x x x	261	emarginatum Prime	x47x
228	Hildrethiana Lea	x x x x3x	262	flavum Prime	x47x
229	marginata Say	x x x x x	263	partumeium Say	x x x x
230	rugosa Bar.	x x x x	264	Jayanum Prime	x47x47x50
231	undulata Say* 43	x	265	sphaericum Anth.	x
232	Anodonta Benedictii Lea	x x2 x x	266	transversum Say	x3x
233	Buchanensis Lea* 44	x2	267	securis Prime	x3x
234	corpulenta Cpr.	x3x		<i>securis crocea</i> Lewis	x
235	decora Lea		268	truncatum Lind.	x x
	<i>inornata</i> Anth.	x x	269	Vermontanum Prime	x
236	edentula Say	x x x x x	270	rosaceum Prime	x x
	<i>edentula rhombica</i> Anth.	x x x	271	stamineum Con.	x x
237	ferruginea Lea	x3	272	tenuis Prime	x
238	Ferussaciana Lea	x x x x x	273	<i>Pisidium Virginicum</i> Bgt.	x x x x
239	fluviatilis Dillw* 45	x x3		<i>dubium</i> Say	x
				<i>dubiosa</i> Say	x

(PAGE 21)

		S	M	C	D	W	
	Anodonta <i>cataracta</i> Say	x	x	x3			
240	Footiana Lea	x	x2x	x			274
	<i>McNeillii</i> Anth.	x	x x				Adamsi Prime 48
	Footiana opalina Anth.	x	x				275
241	fragilis Lam.	x3x					compressum Prime
	<i>flava</i> Anth.	x					276
	<i>glandulosa</i> Anth.	x x					abditum Hald.
	<i>imbricata</i> Anth.	x x					abditum abyssorum Stimp Mss* 49
	<i>irisans</i> Anth.	x x x					277
	<i>pallida</i> Anth.	x x x					rotundatum Prime
	<i>subcarinata</i> Currier	x x x					278
242	imbecilis Say	x x x x					variable Prime
243	lacustris Lea* 45	x3					279
244	Marryatana Lea	x3x					ventricosum Prime
	<i>Houghtonensis</i> Currier	x x x					280
	<i>subinflata</i> Anth.	x x					noveboracense Prime
245	modesta Lea	x x x x					281
246	ovata Lea	x x x x					aequilaterale Prime
	<i>subangulata</i> Anth.	x x					282
							Idahoense Roper
							283
							milium Held
							284
							punctatum Sterki

(PAGE 22)

NOTES.

1. This list is intended to give every species, which has at any time been quoted in any of the formal lists of the state, and such scattering citations of other species as have been found elsewhere, with the name, synonym-

ous or otherwise, given in the original citation. The accepted nomenclature is printed in plain type, synonyms in italics, species considered doubtful are asterisked. The four columns headed respectively S.M.C. and D., contain the species given in the catalogues of Sager, Miles, Currier and DeCamp. In the DeCamp column are included certain species included in my catalogue of 1892, upon the authority of a supplemental written list furnished by Dr. DeCamp for that purpose. In the Currier column certain species are included, given in his 'Catalogue of Grand Rapids Shells,' but not included in his general catalogue of 1868, probably because he became satisfied that the citation was erroneous or synonymous. In the fifth column, headed W, are included all species vouched for by myself, either in my former lists or in the present one. Unless otherwise specified, these citations are based upon specimens in my collection.

2. Cited in 'Grand Rapids' catalogue of 1865 only.

3. Cited in supplemental list only.

4. See Nautilus VII, p. 126, for the circumstances under which this species was found.

5. Has not been found by any recent collector and for the present must be considered doubtful.

6. Leach's name was never published but is cited by Pilsby, Manuel IX, p. 76.

7. The specimens thus cited have been recently described by Sterki as a new species under the name of *V. Morosi*, Nautilus VIII p. 89.

8. Undoubtedly erroneous as the species is confined to the southern region.

9. It is doubtful whether this identification is correct, although specimens have been received under this name said to have been identified by Dr. James Lewis. The form thus designated is a well marked one and seems to be generally distributed through the northern part of the state.

10. The scalariform variety was originally cited by me under this name by mistake.

11. Never described. Said by Dr. DeCamp to be a form of *L. palustris* Mull.

12. Said by Dr. DeCamp to have been found in a greenhouse. It is a western species.

13. Probably erroneous as the species is a western one. A specimen received from Dr. DeCamp under this name is a form of *L. ampla* Migh.

14. Probably erroneous. A purely western species.

15. Cited by Clescin, *Limnaeidae* p. 287.

16. Very doubtful.

17. Has not yet been published. From a drawing kindly furnished by Mr. Beecher, it appears to be closely related to the European *P. nautilus* L.

18. Undoubtedly erroneous as Say's species is Mexican. Miles received his specimen from Currier (Cat. p. 238) and the form is probably that subsequently described as *V. striata* Lewis.

19. Very doubtful, as it has never been found by any of the local collectors. But it may occur along the southern border of the state. Cited only by Haldeman and Binney.

20. Doubtful. Has never been found by any of the recent collectors.

21. Cited upon the authority of Dr. M. L. Leach, who states that a single specimen from Higgins' Lake, Roscommon County, was so identified by Tryon. I think it must be considered as doubtful until its occurrence in Michigan is verified.

22. Cited by Dr. DeCamp, who states that his specimens were identified by

(PAGE 23)

Tryon. The species is a southern one and has not, I believe, been found in any intermediate locality.

23. The species is a Californian one and the identification must be considered doubtful. Specimens received from Dr. DeCamp evidently belong to a different species from any other found in the state and it may be a new one.

24. Clearly erroneous as the species is confined to the Atlantic drainage. As *Pleurocera subulata* Lea is not cited by either Sager or Miles, it seems probable that that is the species referred to.

25. Doubtful. A specimen received from Dr. DeCamp is a rayed *U. rectus* Lam. Cited also by Call (Geo. Cat.), but he informs me that his citation was based solely on information received from Dr. DeCamp.

26. Undoubtedly erroneous as the species is purely a southern one.

27. This species is confined to the Atlantic drainage. The specimens were probably some form of *U. pectens* Lea.

28. Doubtful. A Tennessee species.

29. Monography of the Unionidae p. 60.

30. Cited also by Call. (loc. cit.) *U. novi-eboraci* Lea?
31. Cited by Call (loc. cit.)
32. Doubtful. See Nautilus VI, p. 44.
33. Doubtful. See Nautilus VI, p. 44. The specimen from my own collection there mentioned has been referred to *U. coccineus* Hild. by Mr. C. T. Simpson.
34. Very doubtful. A southern species.
35. Doubtful. Has not been found by any of the recent collectors. *U. Rangianus* Lea may have been the species intended.
36. Cited under an erroneous identification.
37. This species in the United States is confined to the Atlantic drainage. It has, however, been cited from the shore of Lake Superior by Gould and from Manitoba by Christy (J. of C. IV, p. 344) and may be found in the Upper Peninsula. But it must be cited as doubtful until its occurrence is clearly proved. *Unio distans* Anth., although referred to this species, is more likely to be a form of *U. luteolus* Lam.
38. Monography of the Unionidae, p. 53.
39. Doubtful. Specimens received under that name prove to be *U. ventricosus* Bar.
40. Not found by any recent collector. *U. ventricosus* Bar.?
41. Monography of the Unionidae, p. 51.
42. Specimens received from Dr. DeCamp under this name prove to be a small form of *U. luteolus* Lam.
43. Undoubtedly erroneous as the species is peculiar to the Atlantic drainage. As *M. rugosa* Bar. is not cited by Sager, that is probably the species intended.
44. As this species was not included by Currier in his later catalogue, it is probably a mistake.
45. Very doubtful as the species is an eastern one, probably specimens of *A. fragilis* Lam. were in view.
46. Doubtful. See Nautilus VI, p. 65.
47. Cited from 'Lake Superior' probably from original description of Prime. The original locality of *S. flavum*, however, is given as Sault Ste. Marie.
48. Specimens from Holly are cited by Prime in his original description.
49. Not described. Quoted by Smith in his 'Sketch of the Invertebrate Fauna of Lake Superior'.
50. The citation of this species in my catalogue of 1879 from Houghton Lake is probably erroneous.

51. The citation of this species in my former lists was an error, the specimens having proved to be a form of *S. simile* Say.

(PAGE 24)

LITERATURE.

- ADAMS, H. 1861. Proc. Zoöl Soc. London p. 145.
- ANONYMOUS, 1891. Western Range of *Bythinia tentaculata*. Nautilus V, p. 71.
- ANTHONY, J.G., 1865. Descriptions of New Species of N. A. Unionidae. Am. Jour. of Con. I, p. 155.
- ANTHONY, J.G. 1866. Descriptions of New American Fresh-water Shells. Am Jour. of Con. II, p 144.
- BINNEY, Amos. 1851-7. The Terrestrial Air-Breathing Mollusks of the United States, Vol. I, II and III.
- BINNEY, W.G., 1859. The Terrestrial Air-Breathing Mollusks of the United States, Vol. IV.
- BINNEY, W.G., 1878. The Terrestrial Air-Breathing Mollusks of the United States, Vol. V. -Bul. Mus. Comp. Zoöl. Vol. IV
- BINNEY & BLAND. 1869. Land and Fresh-Water Shells of North America. Pt. I. Pulmonata Geophila. -Smith. Misc. Coll. No. 194.
- BINNEY, W.G. 1865. Land and Fresh-Water Shells of North America Pt. II. Pulmonata Limnophila and Thalassophila. -Smith Misc. Coll. No. 143.
- BINNEY, W.G., 1865. Land and Fresh-water Shells of North America. Pt. III. Operculata. -Smith. Misc. Coll. No. 144.
- BINNEY, W.G., 1885. A Manual of American Land Shells. -Bul. U.S. Nat. Mus. No. 23.
- CALL, R. ELLSWORTH. 1885. A Geographic Catalogue of the Unionidae of the Mississippi Valley. Bul. DesMoines Acad. of Sci. Vol. I, p. 5.
- CALL, R. ELLSWORTH. 1886. On the Genus *Campeoloma* Raf. Bul. Wash. Coll. Lab. of N.H. Vol. I, p 149.
- CALL, R. ELLSWORTH. 1894. On the Geographic and Hypsometric Distribution of N.A. Viviparidae. Am Jour. of Sci Vol. XLVIII, p. 132.
- CASE, WILLIAM. 1847. Descriptions of Two New Species of Shells. (with note by A. A. G.) Am. Jour. of Sci. III, p. 101.
- CLESSIN, S. 1879. Die Familie der Cycladeen. Conch. Cab. Ed. 2.
- CLESSIN, S. 1882. Die Familie der Ancylinen. Conch. Cab Ed. 2.

- CONRAD, T.A. 1836. Monography of the Family Unionidae.
 CONRAD, T.A. 1853. Omissions and Corrections to the Synopsis of N.A. Naiades. Proc. Acad. of Nat. Sci. Phila., Vol. VI, p. 449.

(PAGE 25)

- CURRIER, A.O. 1859. List of Shells collected in the Grand River (Mich.) Valley.
 CURRIER, A.O. 1865. Catalogue of the Mollusca of Grand Rapids, Mich. Am Jour. of Con. I, p. 292.
 CURRIER, A.O. 1867. Descriptions of New Species of Fresh-water Shells from Michigan. Am. Jour. of Con. III, p. 112.
 CURRIER, A.O. 1868. List of the Shell-bearing Mollusca of Michigan. Kent Sci. Inst. Misc. Pub. No. 1.
 DeCAMP, W.H. 1881. List of the Shell-bearing Mollusca of Michigan. Kent Sci. Inst. Misc. Pub. No. 5.
 GOULD, A.A. 1847. Notes on Planorbis multivolvis Case, etc. Proc. Bos. Soc. of Nat. Hist. II, pp. 259 and 263.
 GOULD, A.A. 1850. Catalogue of Shells, etc. Lake Superior by L. Agassiz, p. 243.
 HALDEMAN, S.S. 1842. A Monograph of the F.W. Univalve Mollusca of the United States.
 JACKSON, C. T. 1844. List of Shells found at Lake Superior. Proc. Bos. Soc. of Nat. Hist., I, p. 198.
 JAY, John C. 1852. A Catalogue of Shells contained in the Coll. of John C. Jay, 4th Ed.
 KUSTER & CLESSIN. 1838 and 1876. Die Gattung: Anodonta. Conch. Cab. Ed. 2.
 KUSTER, H.C. 1848. Die Flussperlmuscheln: Unio, etc. Conch. Cab. Ed. 2.
 KUSTER, H.C. 1852. Die Gattung: Paludina. Conch. Cab. Ed. 2.
 KUSTER & CLESSIN. 1862 and 1886. Die Familie der Limnaeiden. Conch. Cab. Ed. 2.
 LEA, Isaac. 1830-1874. Observations on the Genus Unio, etc. Reprinted from Trans. Am. Phil. Soc. and Jour. Acad. Nat. Sci. Phila. Vol. I to XIII.
 LEA, Isaac. 1836-70. A synopsis of the Family Unionidae. Four editions.
 LEWIS, James. 1868. On the Shells of Michigan described by A. O. Currier in Vol. III, p. 112, Am. Jour. of Con. Am. Jour. of Con. IV, p. 81.
 LEWIS, James. 1869-70. Observations on Melantho. Am. Jour. of Con. V, p. 33.

- MILES, Manly. 1860. Catalogue of the Mollusks of Michigan. Rep. Geo. Sur. of Mich., 1860, p. 235.
 PILSBRY, H. A. 1893. Manual of Conchology. Second Series, IX, p. 76.
 PRIME, Temple. 1865. Monograph of American Corbiculidae. Smith, Misc. Coll. No. 145.
 ROPER, Edward W. Notes on Sphaerium secure Prime. Nautilus IV, p. 39.

(PAGE 26)

- SAGER, Abram. 1839. Report of the Zoologist of the Geol. Survey of Mich. Doc. H. of Rep. 1839, p. 410. From the pagination given by Binney (Bibliography p. 56), this paper appears to have been also reprinted separately.
 SCUDDER, N.P. 1885. The published writings of Isaac Lea, L.L.D. Bul. U. S. Nat. Mus. No. 23.
 SIMPSON, Chas. T. 1893. On some Fossil Unios and other Fresh-water Shells from the Drift at Toronto, Canada. Proc. U. S. Nat. Mus. XVI, p. 591.
 SMITH, Frank. 1894. List of the Mollusca observed in Lake St. Clair in the Summer of 1893. Bul. Mich. Fish Com No. 4, p. 42.
 SMITH, Sidney I. 1871. Preliminary Report on Dredgings in Lake Superior. Rep. Sec. of War 1871, Pt. 2, p. 1020.
 SMITH, Sidney I. 1872-73. Sketch of the Invertebrate Fauna of Lake Superior. Rep. U. S. Fish Com. 1872-73, Pt. II, p. 690.
 SOWERBY, G. B. Conch. Icon. XX, p. 72.
 STERKI, V. 1894. Vertigo Morsei, n. sp. Nautilus VIII, p. 80.
 STIMPSON, William. 1865. Researches upon the Hydrobinae. Smith, Misc. Coll. No. 201.
 TROMBLEY, Jerome. 1886. Note on Unio Leibii Lea. The Con. Exchange I, p. 26.
 TRYON, G.W. Jr. 1865. Catalogue of the Species of Physa Inhabiting the U. S. Am. Jour. of Con. I, p. 165.
 TRYON, G.W. Jr. 1865. Descriptions of the New Species of N. A. Limnaeidae. Am. Jour. of Con. I, p. 223.
 TRYON, G. W. Jr. 1865. Catalogue of Species of Limnaea Inhabiting the U. S. Am. Jour. of Con. I, p. 247.
 TRYON, G.W. Jr. 1865-6. Monograph of the Family Strepomatidae. Am. Jour. of Con. I, p. 299; II, pp. 14 and 115.

- TRYON, G. W. Jr. 1866. Descriptions of New Fresh-water Shells of the U. S. Am. Jour. of Con. II, p. 4.
- TRYON, G. W. Jr. 1866. Observations on an Abnormal Specimen of *Physa gyrina* Say. Am. Jour. of Con. II, p. 4.
- TRYON, G. W. Jr. 1866-7-8. Monograph of the Terrestrial Mollusca of the U.S. Am. Jour. of Con. II, pp. 218 and 306; III pp. 84, 155 and 298; IV, p. 5. Also printed separately in one volume.
- TRYON, G. W. Jr. 1870. A monograph of the Fresh-water Univalve Mollusca of the United States. A continuation of Haldeman's Monograph of 1842.
- TRYON, G. W. Jr. 1873. Land and Fresh-water Shells of North America. Pt IV: Strepomatidae. = Smiths, Misc Coll. No. 253.
- WALKER & BEECHER 1876. List of Land and Fresh-water Shells found about Ann Arbor, Michigan. Proc. A.A. Sci Assn. for 1876, p. 43.
- (PAGE 27)
- WALKER, Bryant. 1879. Catalogue of the Shell-bearing Mollusca of Michigan. Jour. of Con. II, p. 325.
- WALKER, Bryant. 1888. On the re-discovery of *Planorbis multivolvis* Case and *P. truncatus* Miles. Jour. of Con V, p. 330.
- WALKER, Bryant. 1889. On the Occurrence of *Unio complanatus* Sol. in Michigan. Nautilus III, p. 16.
- WALKER, Bryant. 1891. *Unio complanatus* Sol. in Northern Michigan. Nautilus V, p. 95.
- WALKER, Bryant. 1892. The Shell-bearing Mollusca of Michigan. Nautilus VI, pp. 13, 31, 42, 63 and 135.
- WALKER, Bryant. 1892. Notes on *Unio luteolus* Lam. Nautilus VI, p. 94.
- WALKER, Bryant. 1893. Shells of the Saginaw Valley, Michigan. Nautilus VII, p. 125.

THE TERRESTRIAL MOLLUSCA OF MICHIGAN

BY BRYANT WALKER

Read before the Michigan Academy of Science,
March, 1899.

Printed for the author

DETROIT, MICH.

1899

(PAGE 2)

(M A P)

(PAGE 3)

THE TERRESTRIAL MOLLUSCA OF MICHIGAN.

This paper is based mainly upon the records accumulated by the Conchological Section of the Academy. All other available sources of information, however, have been utilized, and it therefore represents substantially everything that is known at the present time in regard to the extent and distribution of this portion of the fauna of the state.

In the catalogue of the mollusca of the state, which was presented at the first meeting